

## Product datasheet

info@arigobio.com

# ARG10018 anti-G-CSF Receptor antibody [S1390]

Package: 100 μg Store at: -20°C

#### **Summary**

Product Description Mouse Monoclonal antibody [S1390] recognizes G-CSF Receptor

Tested Reactivity Hu
Tested Application WB

Host Mouse

**Clonality** Monoclonal

Clone S1390

Isotype IgG2b, kappa

Target Name G-CSF Receptor

Species Human

Immunogen Purified recombinant MBP-GCSFR fusion protein

Conjugation Un-conjugated

Alternate Names G-CSF receptor; CD114; CD antigen CD114; G-CSF-R; GCSFR; Granulocyte colony-stimulating factor

receptor

### **Application Instructions**

**Application Note** 

Western Blot: The antibody recognizes denatured G-CSFR in Western blot following SDS-PAGE. Suggest use 200µg of cell lysate or immunoprecipitated protein.

\* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

#### **Properties**

Form Liquid

Purification Protein G affinity purified

Buffer 0.01M PBS (pH 7.2)

Concentration 1 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

#### **Bioinformation**

Database links <u>GeneID: 1441 Human</u>

www.arigobio.com arigo.nuts about antibodies 1/2

#### Swiss-port # Q99062 Human

Gene Symbol CSF3R

Gene Full Name colony stimulating factor 3 receptor (granulocyte)

Background The protein encoded by this gene is the receptor for colony stimulating factor 3, a cytokine that

controls the production, differentiation, and function of granulocytes. The encoded protein, which is a member of the family of cytokine receptors, may also function in some cell surface adhesion or recognition processes. Alternatively spliced transcript variants have been described. Mutations in this gene are a cause of Kostmann syndrome, also known as severe congenital neutropenia. [provided by

RefSeq, Aug 2010]

Function Receptor for granulocyte colony-stimulating factor (CSF3), essential for granulocytic maturation. Plays a

crucial role in the proliferation, differientation and survival of cells along the neutrophilic lineage. In

addition it may function in some adhesion or recognition events at the cell surface. [UniProt]

Research Area Cancer antibody; Developmental Biology antibody; Immune System antibody

Calculated Mw 92 kDa

PTM N-glycosylated.