

Product datasheet

info@arigobio.com

ARG21150 anti-CD3 epsilon antibody [CT-3]

Package: 100 μg Store at: -20°C

Summary

Product Description Mouse Monoclonal antibody [CT-3] recognizes CD3 epsilon

Tested Reactivity Chk

Tested Application FACS, IHC-Fr, IHC-P, IP

Specificity Chicken CD3 epsilon. The clone CT-3 recognizes a complex of at least three polypeptides of Mr 20, 19,

and 17 kDa (two of which are N-glycosylated) on chicken T cells. The antibody also coprecipitates a polypeptide of 90 kDa from digitonin solubilized T cell lysates, which can be reduced to two

polypeptides of Mr 50 and 40 kDa.

Host Mouse

Clonality Monoclonal

Clone CT-3

Isotype IgG1, kappa

Target Name CD3 epsilon

Species Chicken

Immunogen Chicken thymocytes and Ig- blood mononuclear cells

Conjugation Un-conjugated

Alternate Names CD3E; CD3 Epsilon Subunit Of T-Cell Receptor Complex; T-Cell Surface Glycoprotein CD3 Epsilon Chain;

CD3e Antigen, Epsilon Polypeptide (TiT3 Complex); T-Cell Surface Antigen T3/Leu-4 Epsilon Chain; CD3e Molecule, Epsilon (CD3-TCR Complex); CD3-Epsilon; CD3epsilon; T3E; T-Cell Antigen Receptor Complex,

Epsilon Subunit Of T3; CD3e Molecule; CD3e Antigen; CD3-EPSILON; CD3EPSILON

Application Instructions

Application table	Application	Dilution
	FACS	Assay-dependent
	IHC-Fr	Assay-dependent
	IHC-P	Assay-dependent
	IP	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Buffer	BBS (pH 8.2)
Concentration	0.5 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol CD3E

Gene Full Name CD3 Epsilon Subunit Of T-Cell Receptor Complex

Background The protein encoded by this gene is the CD3-epsilon polypeptide, which together with CD3-gamma,

-delta and -zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T-cell receptor-CD3 complex. This complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in the same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-cell development. Defects in this gene cause immunodeficiency. This gene has also

been linked to a susceptibility to type I diabetes in women.

Function Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in

adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein

tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathway.

Highlight Related products:

CD3 antibodies; CD3 ELISA Kits; CD3 Duos / Panels;

Related news:

New antibody panels and duos for Tumor immune microenvironment

Tumor-Infiltrating Lymphocytes (TILs)

Research Area Cancer antibody; Developmental Biology antibody; Immune System antibody; Lymphocyte Marker

antibody; Inflammatory Cell Marker antibody; T-cell Marker antibody; T-cell infiltration Study antibody;


Tumor-infiltrating Lymphocyte Study antibody


Calculated Mw 23 kDa

PTM Disulfide bond, Phosphoprotein

Cellular Localization Cell membrane, Membrane

Images





ARG21150 anti-CD3 epsilon antibody [CT-3] IHC-P image

Immunohistochemistry: Paraffin-embedded Chicken spleen tissue stained with ARG21150 anti-CD3 epsilon antibody [CT-3] (right) followed by anti-Mouse IgG antibody (HRP), DAB and hematoxylin.