

ARG21487 anti-IL17 / IL17A antibody [TC11-8H4] (FITC)

Package: 50 µg
Store at: 4°C

Summary

Product Description	FITC-conjugated Rat Monoclonal antibody [TC11-8H4] recognizes IL17 / IL17A
Tested Reactivity	Ms
Tested Application	ELISA, ELISPOT, FACS
Specificity	Mouse IL-17A.
Host	Rat
Clonality	Monoclonal
Clone	TC11-8H4
Isotype	IgG1, kappa
Target Name	IL17 / IL17A
Species	Mouse
Immunogen	E. coli-expressed mouse IL-17
Conjugation	FITC
Alternate Names	IL-17A; Interleukin-17A; IL-17; CTLA8; IL17; CTLA-8; Cytotoxic T-lymphocyte-associated antigen 8

Application Instructions


Application table	Application	Dilution
	ELISA	Assay-dependent
	ELISPOT	Assay-dependent
	FACS	< 1 µg/10 ⁶ cells
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Buffer	PBS and 0.1% Sodium azide.
Preservative	0.1% Sodium azide
Concentration	0.1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Database links	GeneID: 16171 Mouse Swiss-port # Q62386 Mouse
Gene Symbol	IL17A
Gene Full Name	interleukin 17A
Background	The protein encoded by this gene is a proinflammatory cytokine produced by activated T cells. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis. [provided by RefSeq, Jul 2008]
Function	Induces stromal cells to produce proinflammatory and hematopoietic cytokines. Enhances the surface expression of ICAM1/intracellular adhesion molecule 1 in fibroblasts. [UniProt]
Highlight	Related products: IL17 antibodies ; IL17 ELISA Kits ; IL17 recombinant proteins ; Anti-Rat IgG secondary antibodies ; Related news: HMGB1, a biomarker and therapeutic target in COVID-19
Calculated Mw	18 kDa
PTM	Found both in glycosylated and nonglycosylated forms.

Images



ARG21487 anti-IL17 / IL17A antibody [TC11-8H4] (FITC) FACS image

Flow Cytometry: PMA and Ionomycin-stimulated C57BL/6N Mouse T lymphocyte EL-4 cell line stained with ARG21487 anti-IL17 / IL17A antibody [TC11-8H4] (FITC).