

ARG21693 Goat anti-Rat IgG antibody (PE), pre-adsorbed

Package: 250 µg
Store at: 4°C

Summary

Product Description	PE-conjugated Goat Polyclonal antibody recognizes Rat IgG
Tested Reactivity	Rat
Tested Application	FACS, FLISA, ICC/IF, IHC-Fr, WB
Specificity	The antibody reacts with the heavy chain of Rat IgG. The antibody is pre-adsorbed with Rat IgM, so the antibody may not react with Rat IgM, but may react with immunoglobulins from other species.
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	IgG
Species	Rat
Conjugation	PE

Application Instructions

Pre Adsorbed	Rat IgM.												
Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>FACS</td><td>Assay-dependent</td></tr><tr><td>FLISA</td><td>< 1 ug/ml</td></tr><tr><td>ICC/IF</td><td>Assay-dependent</td></tr><tr><td>IHC-Fr</td><td>Assay-dependent</td></tr><tr><td>WB</td><td>Assay-dependent</td></tr></tbody></table>	Application	Dilution	FACS	Assay-dependent	FLISA	< 1 ug/ml	ICC/IF	Assay-dependent	IHC-Fr	Assay-dependent	WB	Assay-dependent
Application	Dilution												
FACS	Assay-dependent												
FLISA	< 1 ug/ml												
ICC/IF	Assay-dependent												
IHC-Fr	Assay-dependent												
WB	Assay-dependent												
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.												

Properties


Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS, 0.1% Sodium azide and Sucrose.
Preservative	0.1% Sodium azide
Stabilizer	Sucrose
Concentration	0.5 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be

gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

Images



ARG21693 Goat anti-Rat IgG antibody (PE) (pre-adsorbed) FLISA image

FLISA: The plate was coated with purified Rat IgG and IgM. Immunoglobulins were detected with serially diluted ARG21693 Goat anti-Rat IgG antibody (PE) (pre-adsorbed).