

ARG21717 Goat anti-Rabbit IgM antibody, pre-adsorbed

Package: 500 µg
Store at: -20°C

Summary

| | |
|---------------------|---|
| Product Description | Goat Polyclonal antibody recognizes Rabbit IgM |
| Tested Reactivity | Rb |
| Tested Application | ELISA, ELISPOT, FACS, FLISA, IHC-Fr, IHC-P, WB |
| Specificity | The antibody reacts with the heavy chain of Rabbit IgM. The antibody is pre-adsorbed with Rabbit IgG, so the antibody may not react with Rabbit IgG, but may react with immunoglobulins from other species. |
| Host | Goat |
| Clonality | Polyclonal |
| Isotype | IgG |
| Target Name | IgM |
| Species | Rabbit |
| Conjugation | Un-conjugated |

Application Instructions

| Pre Adsorbed | Rabbit IgG. | | | | | | | | | | | | | | | | |
|-------------------|---|-------------|----------|-------|-----------------|---------|-----------------|------|-----------------|-------|-----------------|--------|-----------------|-------|-----------------|----|-----------------|
| Application table | <table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>ELISA</td><td>Assay-dependent</td></tr><tr><td>ELISPOT</td><td>Assay-dependent</td></tr><tr><td>FACS</td><td>Assay-dependent</td></tr><tr><td>FLISA</td><td>Assay-dependent</td></tr><tr><td>IHC-Fr</td><td>Assay-dependent</td></tr><tr><td>IHC-P</td><td>Assay-dependent</td></tr><tr><td>WB</td><td>Assay-dependent</td></tr></tbody></table> | Application | Dilution | ELISA | Assay-dependent | ELISPOT | Assay-dependent | FACS | Assay-dependent | FLISA | Assay-dependent | IHC-Fr | Assay-dependent | IHC-P | Assay-dependent | WB | Assay-dependent |
| Application | Dilution | | | | | | | | | | | | | | | | |
| ELISA | Assay-dependent | | | | | | | | | | | | | | | | |
| ELISPOT | Assay-dependent | | | | | | | | | | | | | | | | |
| FACS | Assay-dependent | | | | | | | | | | | | | | | | |
| FLISA | Assay-dependent | | | | | | | | | | | | | | | | |
| IHC-Fr | Assay-dependent | | | | | | | | | | | | | | | | |
| IHC-P | Assay-dependent | | | | | | | | | | | | | | | | |
| WB | Assay-dependent | | | | | | | | | | | | | | | | |
| Application Note | * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. | | | | | | | | | | | | | | | | |

Properties

| | |
|---------------------|--|
| Form | Liquid |
| Purification | Affinity purification with immunogen. |
| Buffer | BBS (pH 8.2) |
| Concentration | 1 mg/ml |
| Storage instruction | For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw |

cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.