

## ARG21915 Goat anti-Human IgA antibody, F(ab')<sub>2</sub> fragment (Biotin), pre-adsorbed

Package: 250 µg  
Store at: 4°C

### Summary

Product Description	Biotin-conjugated F(ab') <sub>2</sub> fragment of Goat Polyclonal antibody recognizes Human IgA
Tested Reactivity	Hu
Tested Application	ELISA, ELISPOT, FACS, FLISA, IHC-Fr
Specificity	The antibody reacts with the heavy chain of Human IgA. The antibody is pre-adsorbed with Human IgG and IgM, so the antibody may not react with Human IgG and IgM, but may react with IgA from other species.
Host	Goat
Clonality	Polyclonal
Isotype	F(ab') <sub>2</sub> IgG
Target Name	IgA
Species	Human
Conjugation	Biotin

### Application Instructions

Pre Adsorbed	Human IgG and IgM.	
Application table	Application	Dilution
	ELISA	1:5000 - 1:20000
	ELISPOT	Assay-dependent
	FACS	< 1 µg/10 <sup>6</sup> cells
	FLISA	Assay-dependent
	IHC-Fr	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Buffer	PBS and 0.1% Sodium azide.
Preservative	0.1% Sodium azide
Concentration	0.5 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be

gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.


## Bioinformation

Highlight

Related news:

[Comprehensive anti-Human secondary antibodies](#)

## Images



ARG21915 Goat anti-Human IgA antibody, F(ab')<sub>2</sub> fragment (Biotin) (pre-adsorbed) ELISA image

ELISA: The plate was coated with purified Human IgA, IgG, and IgM. Immunoglobulins were detected with serially diluted ARG21915 Goat anti-Human IgA antibody, F(ab')<sub>2</sub> fragment (Biotin) (pre-adsorbed) followed by [ARG23912](#) Streptavidin (HRP).