

ARG21925 Goat anti-Bovine IgG (H+L) antibody (Biotin)

Package: 1 mg
Store at: 4°C

Summary

Product Description	Biotin-conjugated Goat Polyclonal antibody recognizes Bovine IgG (H+L)
Tested Reactivity	Bov
Tested Application	ELISA, FACS, FLISA, ICC/IF, IP, Puri, WB
Specificity	Reacts with the heavy and light chains of bovine IgG
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	IgG (H+L)
Species	Bovine
Conjugation	Biotin

Application Instructions

Application table	Application	Dilution
	ELISA	1:5000 - 1:20000
	FACS	Assay-dependent
	FLISA	Assay-dependent
	ICC/IF	Assay-dependent
	IP	Assay-dependent
	Puri	Assay-dependent
	WB	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS and 0.1% Sodium azide.
Preservative	0.1% Sodium azide
Concentration	0.5 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.