

ARG21936 Goat anti-Dog IgG (H+L) antibody (Biotin)

Package: 1 mg
Store at: 4°C

Summary

| | |
|---------------------|---|
| Product Description | Biotin-conjugated Goat Polyclonal antibody recognizes Dog IgG (H+L) |
| Tested Reactivity | Dog |
| Tested Application | ELISA, FLISA, ICC/IF, IHC-Fr, WB |
| Specificity | Reacts with the heavy and light chains of Dog IgG |
| Host | Goat |
| Clonality | Polyclonal |
| Isotype | IgG |
| Target Name | IgG (H+L) |
| Species | Dog |
| Conjugation | Biotin |

Application Instructions

| Application table | Application | Dilution |
|-------------------|-------------|------------------|
| | ELISA | 1:5000 - 1:20000 |
| | FLISA | Assay-dependent |
| | ICC/IF | Assay-dependent |
| | IHC-Fr | Assay-dependent |
| | WB | Assay-dependent |

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

| | |
|---------------------|--|
| Form | Liquid |
| Purification | Affinity purification with immunogen. |
| Buffer | PBS and 0.1% Sodium azide. |
| Preservative | 0.1% Sodium azide |
| Concentration | 0.5 mg/ml |
| Storage instruction | Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |
| Note | For laboratory research only, not for drug, diagnostic or other use. |