

ARG22304 Goat anti-Mouse IgM antibody (HRP), pre-adsorbed

Package: 500 µl
Store at: 4°C

Summary

Product Description	HRP-conjugated Goat Polyclonal antibody recognizes Mouse IgM
Tested Reactivity	Ms
Tested Application	ELISA, ELISPOT, EM, FACS, FLISA, ICC/IF, IHC-Fr, Puri, WB
Specificity	Reacts with the heavy chain of mouse IgM
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	IgM
Species	Mouse
Immunogen	Mouse IgM
Conjugation	HRP

Application Instructions

Pre Adsorbed	Human																				
Application table	<table><tr><th>Application</th><th>Dilution</th></tr><tr><td>ELISA</td><td>1:4000 - 1:8000</td></tr><tr><td>ELISPOT</td><td>Assay-dependent</td></tr><tr><td>EM</td><td>Assay-dependent</td></tr><tr><td>FACS</td><td>Assay-dependent</td></tr><tr><td>FLISA</td><td>Assay-dependent</td></tr><tr><td>ICC/IF</td><td>Assay-dependent</td></tr><tr><td>IHC-Fr</td><td>Assay-dependent</td></tr><tr><td>Puri</td><td>Assay-dependent</td></tr><tr><td>WB</td><td>Assay-dependent</td></tr></table>	Application	Dilution	ELISA	1:4000 - 1:8000	ELISPOT	Assay-dependent	EM	Assay-dependent	FACS	Assay-dependent	FLISA	Assay-dependent	ICC/IF	Assay-dependent	IHC-Fr	Assay-dependent	Puri	Assay-dependent	WB	Assay-dependent
	Application	Dilution																			
	ELISA	1:4000 - 1:8000																			
	ELISPOT	Assay-dependent																			
	EM	Assay-dependent																			
	FACS	Assay-dependent																			
	FLISA	Assay-dependent																			
	ICC/IF	Assay-dependent																			
	IHC-Fr	Assay-dependent																			
	Puri	Assay-dependent																			
WB	Assay-dependent																				
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.																				

Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS (pH 7.4) and 50% Glycerol

Stabilizer	50% Glycerol
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Images

