

Product datasheet

info@arigobio.com

ARG22676 anti-Ly-6B.2 antibody [7/4] (FITC)

Package: 50 μg Store at: 4°C

Summary

Product Description FITC-conjugated Rat Monoclonal antibody [7/4] recognizes Ly-6B.2

This antibody recognizes Ly-6B.2 antigen. Ly-6B.2 is a ~25-30kDa GPI-anchored, heavily glycosylated protein expressed on neutrophils, inflammatory monocytes and some activated macrophages (Rosas et al. 2010). High levels of expression are seen in bone marrow, spleen, lung and lymph nodes. N-glycanase treatment of thioglycollate elicited peritoneal neutrophil lysates lowers the apparent molecular weight of Ly-6B.2 to ~15kDa (Rosas et al.2010).

In common with other Ly-6 antigens Ly-6B.2 demonstrates a polymorphic expression on inbred mouse strains (Kimura et al. 1984). Rat anti mouse Ly-6B.2, clone 7/4 recognizes the Ly-6B.2 antigen in 129J; AKR; C57BL/6; C57BL/10; C58; DBA/2; NZB; NZW; SJL; MFI; Swiss (PO) Strains whilst A2G; A/Sn; ASW; BALB/c; C3H/HEH: CBA.

T6T6 are negative or demonstrate very weak reactivity (Hirsch and Gordon 1982).

Rat anti mouse Ly-6B.2 has been successfully used for the immunomagnetic depletion of neutrophils during the enrichment of primitive hematopoietic cells from bone marrow (Bertoncello et al. 1991) and the depletion of myeloid cells in vivo (Rosas et al. 2010).

Tested Reactivity Ms

Tested Application FACS

Host Rat

Clonality Monoclonal

Clone 7/4

Isotype IgG2a

Target Name Ly-6B.2

Species Mouse

Immunogen Cultured bone marrow cells

Conjugation FITC

Application Instructions

Application table	Application	Dilution
	FACS	Neat - 1:10
Application Note	FACS: Use 10 μ l of the suggested working dilution to label 10^6 cells in 100 μ l.	

st The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations

should be determined by the scientist.

Properties

Form Liquid

Purification Purification with Protein G.

Buffer PBS, 0.09% Sodium azide and 1% BSA.

Preservative 0.09% Sodium azide

Stabilizer 1% BSA

Concentration 0.1 mg/ml

Storage instruction Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid

repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be

gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.