

## ARG22809 anti-CD4 antibody [YKIX302.9] (FITC)

Package: 50 tests  
Store at: 4°C

### Summary

Product Description	FITC-conjugated Rat Monoclonal antibody [YKIX302.9] recognizes CD4 Rat anti Dog CD4 antibody, clone YKIX302.9, is a monoclonal antibody specific for the canine CD4 cell surface antigen. Clone YKIX302.9 was clustered at the first Canine Leukocyte Antigen Workshop (CLAW) [Cobbold et al. 1992] along with clone CA13.1E4. Rat anti Dog CD4 (YKIX302.9) has been demonstrated to partially deplete circulating T lymphocytes when administered in vivo, but alone was not sufficient to prolong allograft survival in a canine transplant model (Watson et al. 1993). Uniquely amongst mammals, canine CD4 is expressed by neutrophils as well as by lymphocytes subsets Moore et al. 1992. Rat anti Canine CD4 (YKIX302.9) forms part of a panel of anti canine monoclonal antibodies used extensively in the evaluation of leukemic status in dogs (Villiers 2002).
Tested Reactivity	Dog
Tested Application	FACS
Host	Rat
Clonality	Monoclonal
Clone	YKIX302.9
Isotype	IgG2a
Target Name	CD4
Species	Dog
Immunogen	Canine concanavilin A activated T cell blasts.
Conjugation	FITC
Alternate Names	CD4mut; CD antigen CD4; T-cell surface glycoprotein CD4; T-cell surface antigen T4/Leu-3

### Application Instructions

Application table	<table> <tr> <th>Application</th><th>Dilution</th></tr> <tr> <td>FACS</td><td>Neat - 1:10</td></tr> </table>	Application	Dilution	FACS	Neat - 1:10
Application	Dilution				
FACS	Neat - 1:10				
Application Note	<p>FACS: Use 10 µl of the suggested working dilution to label 10<sup>6</sup> cells or 100 µl whole blood.</p> <p>* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.</p>				


### Properties

Form	Liquid
Purification	Purification with Protein G.
Buffer	PBS, 0.09% Sodium azide and 1% BSA
Preservative	0.09% Sodium azide
Stabilizer	1% BSA

Concentration	0.1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


## Bioinformation

Gene Symbol	CD4
Gene Full Name	CD4 molecule
Background	CD4 is a membrane glycoprotein of T lymphocytes that interacts with major histocompatibility complex class II antigens and is also a receptor for the human immunodeficiency virus. This gene is expressed not only in T lymphocytes, but also in B cells, macrophages, and granulocytes. It is also expressed in specific regions of the brain. The protein functions to initiate or augment the early phase of T-cell activation, and may function as an important mediator of indirect neuronal damage in infectious and immune-mediated diseases of the central nervous system. Multiple alternatively spliced transcript variants encoding different isoforms have been identified in this gene. [provided by RefSeq, Aug 2010]
Function	CD4 is an integral membrane glycoprotein that plays an essential role in the immune response and serves multiple functions in responses against both external and internal offenses. In T-cells, functions primarily as a coreceptor for MHC class II molecule:peptide complex. The antigens presented by class II peptides are derived from extracellular proteins while class I peptides are derived from cytosolic proteins. Interacts simultaneously with the T-cell receptor (TCR) and the MHC class II presented by antigen presenting cells (APCs). In turn, recruits the Src kinase LCK to the vicinity of the TCR-CD3 complex. LCK then initiates different intracellular signaling pathways by phosphorylating various substrates ultimately leading to lymphokine production, motility, adhesion and activation of T-helper cells. In other cells such as macrophages or NK cells, plays a role in differentiation/activation, cytokine expression and cell migration in a TCR/LCK-independent pathway. Participates in the development of T-helper cells in the thymus and triggers the differentiation of monocytes into functional mature macrophages. [UniProt]
Highlight	Related products: <a href="#">CD4 antibodies</a> ; <a href="#">CD4 ELISA Kits</a> ; <a href="#">CD4 Duos / Panels</a> ; <a href="#">Anti-Rat IgG secondary antibodies</a> ; Related news: <a href="#">New antibody panels and duos for Tumor immune microenvironment</a> <a href="#">Tumor-Infiltrating Lymphocytes (TILs)</a>
Research Area	Developmental Biology antibody; Immune System antibody; Regulatory T cells Study antibody; T-cell infiltration Study antibody; Tumor-infiltrating Lymphocyte Study antibody
Calculated Mw	51 kDa
PTM	Palmitoylation and association with LCK contribute to the enrichment of CD4 in lipid rafts.



ARG22809 anti-CD4 antibody [YKIX302.9] (FITC) FACS image

Flow Cytometry: Canine peripheral blood lymphocytes stained with ARG22809 anti-CD4 antibody [YKIX302.9] (FITC).



ARG22809 anti-CD4 antibody [YKIX302.9] (FITC) FACS image

Flow Cytometry: Figure A. Purified Mouse anti Canine CD8b detected with Goat anti Mouse IgG1 DyLight 649 and Rat IgG2b FITC isotype control. Figure B. Purified Mouse anti Canine CD8b detected with Goat anti Mouse IgG1 PE and ARG22809 anti-CD4 antibody [YKIX302.9] (FITC). All experiments performed on red cell lysed canine blood gated on mononuclear cells.