

# Product datasheet

info@arigobio.com

# ARG23494 anti-Albumin antibody [BGN/D1]

Package: 100 μg Store at: -20°C

#### **Summary**

Product Description Mouse Monoclonal antibody [BGN/D1] recognizes Albumin.

Rabbit anti Bovine serum albumin antibody recognizes bovine albumin, a major protein constituant of

bovine serum.

Tested Reactivity Bov

Species Does Not React With Hu, Ms, Rat, Goat, Hrs, Rb

Tested Application ELISA, WB

Host Mouse

Clonality Monoclonal
Clone BGN/D1

Isotype IgG1

Target Name Albumin
Species Bovine

Immunogen Bovine serum albumin.

Conjugation Un-conjugated

Alternate Names FDAH; PRO0883; PRO1341; Serum albumin; ANALBA; PRO0903

## **Application Instructions**

Application table	Application	Dilution
	ELISA	1:250 - 1:1000
	WB	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### **Properties**

Form Liquid

Purification Purification with Protein G.

Buffer PBS and 0.09% Sodium azide.

Preservative 0.09% Sodium azide

Concentration 1 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

#### Bioinformation

Gene Symbol ALB

Gene Full Name albumin

Background Albumin is a soluble, monomeric protein which comprises about one-half of the blood serum protein.

Albumin functions primarily as a carrier protein for steroids, fatty acids, and thyroid hormones and plays a role in stabilizing extracellular fluid volume. Albumin is a globular unglycosylated serum protein of molecular weight 65,000. Albumin is synthesized in the liver as preproalbumin which has an N-terminal peptide that is removed before the nascent protein is released from the rough endoplasmic reticulum. The product, proalbumin, is in turn cleaved in the Golgi vesicles to produce the secreted

albumin. [provided by RefSeq, Jul 2008]

Function Serum albumin, the main protein of plasma, has a good binding capacity for water, Ca(2+), Na(+), K(+),

fatty acids, hormones, bilirubin and drugs. Its main function is the regulation of the colloidal osmotic pressure of blood. Major zinc transporter in plasma, typically binds about 80% of all plasma zinc.

[UniProt]

Calculated Mw 69 kDa

PTM Kenitra variant is partially O-glycosylated at Thr-620. It has two new disulfide bonds Cys-600 to Cys-602

and Cys-601 to Cys-606.

Glycated in diabetic patients.

Phosphorylated by FAM20C in the extracellular medium.

Acetylated on Lys-223 by acetylsalicylic acid. [UniProt]