

## ARG30334 Tumor-infiltrating Lymphocyte Antibody Panel

Package: 1 kit  
Store at: -20°C

### Component

Cat. No.	Component Name	Host clonality	Reactivity	Application	Package
ARG65859	anti-CD3 epsilon antibody [SQab1713]	Rabbit mAb	Hu	FACS, ICC/IF, IHC-P, IP, WB	20 µl
ARG66628	anti-CD8 antibody [SQab19146]	Rabbit mAb	Hu	IHC-P	20 µl
ARG65860	anti-CD4 antibody [SQab1714]	Rabbit mAb	Hu	FACS, IHC-P, IP, WB	20 µl
ARG66197	anti-CD20 antibody [SQab1719]	Rabbit mAb	Hu	FACS, ICC/IF, IHC-P, IP	20 µl

### Summary

**Product Description** Tumor-infiltrating Lymphocyte (TIL) Antibody Panel is an all-in-one solution to make identification of TILs easy and economic. This antibody panel comprises antibodies against CD3, CD4, CD8 and CD20 for identifying general T cells, Treg cells, cytotoxic T cells and B cells, respectively. All the antibodies in this panel have excellent IHC staining performance.

Related news:

[New antibody panels and duos for Tumor immune microenvironment Tumor-Infiltrating Lymphocytes \(TILs\)](#)

**Target Name** Tumor-infiltrating Lymphocyte

**Alternate Names** Tumor-infiltrating Lymphocyte antibody; CD3 antibody; CD4 antibody; CD20 antibody; CD8 antibody

### Properties

**Storage instruction** For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

**Note** For laboratory research only, not for drug, diagnostic or other use.


### Bioinformation

**Gene Full Name** Antibody Panel for Tumor-infiltrating Lymphocyte

**Highlight** Related Product:  
[anti-CD3 epsilon antibody;](#)  
[anti-CD8 antibody;](#)  
[anti-CD4 antibody;](#)  
[anti-CD20 antibody;](#)


**Research Area** Tumor infiltrating Lymphocyte markers; Immunology; anti-tumor immune responses; tumor immune

## Images




ARG65859 anti-CD3 epsilon antibody [SQab1713] WB image (Customer's Feedback)

Western blot: 20 µg of Jurkat and Mouse spleen (untreated or treated with LPS) lysates stained with ARG65859 anti-CD3 epsilon antibody [SQab1713] at 1:1000 dilution, overnight at 4°C.




ARG65859 anti-CD3 epsilon antibody [SQab1713] WB image (Customer's Feedback)

Western blot: 30 µg of Molt4 cell lysate stained with ARG65859 anti-CD3 epsilon antibody [SQab1713] at 1:500 dilution.




ARG65859 anti-CD3 epsilon antibody [SQab1713] ICC/IF image

Immunofluorescence: Jurkat cells were fixed with 4% paraformaldehyde for 30 min at RT, permeabilized with 0.1% Triton X-100 for 10 min at RT then blocked with 10% Goat serum for half an hour at room temperature. Samples were stained with ARG65859 anti-CD3 epsilon antibody [SQab1713] (green) at 1:50 and 4°C. DAPI (blue) was used as the nuclear counter stain. Control: PBS and secondary antibody.




#### ARG65859 anti-CD3 epsilon antibody [SQab1713] FACS image

Flow Cytometry: Jurkat cells were fixed with 4% paraformaldehyde for 10 min. The cells were then stained with ARG65859 anti-CD3 epsilon antibody [SQab1713] (blue) at 1:1000 dilution in 1x PBS/1% BSA for 30 min at room temperature, followed by Alexa Fluor® 488 labelled secondary antibody. Unlabelled sample (red) was used as a control.



#### ARG65859 anti-CD3 epsilon antibody [SQab1713] IHC-P image


Immunohistochemistry: Formalin/PFA-fixed and paraffin-embedded sections of Human tonsil tissue stained with ARG65859 anti-CD3 epsilon antibody [SQab1713] at 1:200 dilution. Antigen Retrieval: Boil tissue section in Tris/EDTA buffer (pH 9.0).



#### ARG65859 anti-CD3 epsilon antibody [SQab1713] IP image


Immunoprecipitation: 0.4 mg of Molt-4 whole cell lysate was immunoprecipitated (1:15 dilution) and stained with ARG65859 anti-CD3 epsilon antibody [SQab1713].

Lane 1: Immunoprecipitation in Molt-4 whole cell lysate  
Lane 2: Rabbit IgG instead of Primary Ab in Molt-4 whole cell lysate  
Lane 3: Molt-4 whole cell lysate, 10 µg (input)



#### ARG65859 anti-CD3 epsilon antibody [SQab1713] IHC-P image


Immunohistochemistry: Formalin/PFA-fixed and paraffin-embedded sections of Human colon tissue stained with ARG65859 anti-CD3 epsilon antibody [SQab1713] at 1:200 dilution. Antigen Retrieval: Boil tissue section in Tris/EDTA buffer (pH 9.0).



#### ARG66628 anti-CD8 antibody [SQab19146] IHC-P image


Immunohistochemistry: Human nonacral melanoma and acral melanoma tumor stained with ARG66628 anti-CD8 antibody [SQab19146], [ARG66243 anti-CD279 / PD-1 antibody \[SQab1732\]](#) and [ARG65862 anti-CD274 / PD-L1 antibody \[SQab1716\]](#).

From Yu-Jen Chiu et al. J Dermatol. (2024), [doi: 10.1111/1346-8138.17187](#), Fig. 7.




#### ARG66628 anti-CD8 antibody [SQab19146] IHC-P image

Immunohistochemistry: Formalin/PFA-fixed and paraffin-embedded Human tonsil tissue stained with ARG66628 anti-CD8 antibody [SQab19146]. Antigen Retrieval: Heat mediation was performed in Tris/EDTA buffer (pH 9.0).




#### ARG65860 anti-CD4 antibody [SQab1714] WB image

Western blot: 30 µg of Jurkat and Molt4 cell lysates stained with ARG65860 anti-CD4 antibody [SQab1714] at 1:500 dilution.




#### ARG65860 anti-CD4 antibody [SQab1714] FACS image

Flow Cytometry: Jurkat cells were fixed with 4% paraformaldehyde for 10 min. The cells were then stained with ARG65860 anti-CD4 antibody [SQab1714] (blue) at 1:50 dilution in 1x PBS/1% BSA for 30 min at room temperature, followed by Alexa Fluor® 488 labelled secondary antibody. Unlabelled sample (red) was used as a control.



ARG65860 anti-CD4 antibody [SQab1714] IHC-P image

Immunohistochemistry: Formalin/PFA-fixed and paraffin-embedded sections of Human tonsil tissue stained with ARG65860 anti-CD4 antibody [SQab1714] at 1:2000 dilution. Antigen Retrieval: Boil tissue section in Tris/EDTA buffer (pH 9.0).




ARG65860 anti-CD4 antibody [SQab1714] IP image

Immunoprecipitation: 0.4 mg of Molt-4 whole cell lysate was immunoprecipitated (1:50 dilution) and stained with ARG65860 anti-CD4 antibody [SQab1714].

Lane 1: Immunoprecipitation in Molt-4 whole cell lysate


Lane 2: Rabbit IgG instead of Primary Ab in Molt-4 whole cell lysate

Lane 3: Molt-4 whole cell lysate, 10 µg (input)



ARG66197 anti-CD20 antibody [SQab1719] ICC/IF image


Immunofluorescence: Raji cells fixed with 4% paraformaldehyde for 30 min at RT, permeabilized with 0.1% Triton X-100 for 10 min at RT then blocked with 10% Goat serum for half an hour at room temperature. Samples were stained with ARG66197 anti-CD20 antibody [SQab1719] (green) at 1:1000 at 4°C. DAPI (blue) was used as the nuclear counter stain. Control: PBS and secondary antibody.



ARG66197 anti-CD20 antibody [SQab1719] FACS image

Flow Cytometry: Raji cells were fixed with 4% paraformaldehyde for 10 min. The cells were then stained with ARG66197 anti-CD20 antibody [SQab1719] (red) at 1:500 dilution in 1x PBS/1% BSA for 30 min at room temperature, followed by Alexa Fluor® 488 labelled secondary antibody. Unlabelled sample (black) was used as a control.






#### ARG66197 anti-CD20 antibody [SQab1719] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human appendix tissue stained with ARG66197 anti-CD20 antibody [SQab1719] at 1:20000 dilution.


Antigen retrieval: Heat mediated was performed using Tris/EDTA buffer pH 9.0



#### ARG66197 anti-CD20 antibody [SQab1719] IP image

Immunoprecipitation: 0.4 mg of Raji whole cell lysate was immunoprecipitated (1:20 dilution) and stained with ARG66197 anti-CD20 antibody [SQab1719].

Lane 1: Immunoprecipitation in Raji whole cell lysate  
Lane 2: PBS instead of Primary Ab in Raji whole cell lysate  
Lane 3: Raji whole cell lysate, 10 µg (input)



#### ARG66197 anti-CD20 antibody [SQab1719] IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human spleen tissue stained with ARG66197 anti-CD20 antibody [SQab1719] at 1:20000 dilution.

Antigen retrieval: Heat mediated was performed using Tris/EDTA buffer pH 9.0