

ARG40208 anti-AMACR / p504S antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes AMACR / p504S
Tested Reactivity	Hu, Rat
Predict Reactivity	Bov, Hrs, Mk
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	AMACR / p504S
Species	Human
Immunogen	Synthetic peptide corresponding to aa. 208-246 of Human AMACR / p504S. (RGQNMLDGGAPFYTTYRTADGEFMAVGAIEPQFYELLK)
Conjugation	Un-conjugated
Alternate Names	AMACR; Alpha-Methylacyl-CoA Racemase; P504S; RACE; 2-Methylacyl-CoA Racemase; EC 5.1.99.4; AMACRD; CBAS4; RM

Application Instructions

Application table	Application	Dilution
	WB	0.1 - 0.5 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.05% Sodium azide and 5% BSA.
Preservative	0.05% Sodium azide
Stabilizer	5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	AMACR
Gene Full Name	alpha-methylacyl-CoA racemase
Background	This gene encodes a racemase. The encoded enzyme interconverts pristanoyl-CoA and C27-bile acylCoAs between their (R)- and (S)-stereoisomers. The conversion to the (S)-stereoisomers is necessary for degradation of these substrates by peroxisomal beta-oxidation. Encoded proteins from this locus localize to both mitochondria and peroxisomes. Mutations in this gene may be associated with adult-onset sensorimotor neuropathy, pigmentary retinopathy, and adrenomyeloneuropathy due to defects in bile acid synthesis. Alternatively spliced transcript variants have been described. Read-through transcription also exists between this gene and the upstream neighboring C1QTNF3 (C1q and tumor necrosis factor related protein 3) gene.
Function	Acts only on coenzyme A thioesters, not on free fatty acids, and accepts as substrates a wide range of alpha-methylacyl-CoAs, including pristanoyl-CoA, trihydroxycoprostanoyl-CoA (an intermediate in bile acid synthesis), and arylpropionic acids like the anti-inflammatory drug ibuprofen (2-(4-isobutylphenyl)propionic acid) but neither 3-methyl-branched nor linear-chain acyl-CoAs.
Calculated Mw	42 kDa
PTM	Acetylation
Cellular Localization	Mitochondrion, Peroxisome

Images



ARG40208 anti-AMACR / p504S antibody WB image

Western blot: Rat kidney, Rat liver and HepG2 whole cell lysates stained with ARG40208 anti-AMACR / p504S antibody at 0.5 µg/ml dilution.