

## ARG40479 anti-MCM6 antibody

Package: 50 µg  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes MCM6
Tested Reactivity	Hu, Ms, Rat
Predict Reactivity	Hm
Tested Application	FACS, ICC/IF, IHC-Fr, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	MCM6
Species	Human
Immunogen	Synthetic peptide corresponding to aa. 589-605 of Human MCM6. (ESEDFIVEQYKHLRQRD)
Conjugation	Un-conjugated
Alternate Names	p105MCM; P105MCM; DNA replication licensing factor MCM6; MCG40308; EC 3.6.4.12; Mis5

### Application Instructions

Application table	Application	Dilution
	FACS	1:150 - 1:500
	ICC/IF	1:200 - 1:1000
	IHC-Fr	1:200 - 1:1000
	IHC-P	0.5 - 1 µg/ml
	WB	0.1 - 0.5 µg/ml
Application Note	IHC-P: Antigen Retrieval: By heat mediation. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties


Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.2% Na <sub>2</sub> HPO <sub>4</sub> , 0.9% NaCl, 0.05% Thimerosal, 0.05% Sodium azide and 5% BSA.
Preservative	0.05% Thimerosal and 0.05% Sodium azide
Stabilizer	5% BSA
Concentration	0.5 mg/ml

Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


Gene Symbol	MCM6
Gene Full Name	minichromosome maintenance complex component 6
Background	The protein encoded by this gene is one of the highly conserved mini-chromosome maintenance proteins (MCM) that are essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex (pre_RC) and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 7 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. The phosphorylation of the complex by CDC2 kinase reduces the helicase activity, suggesting a role in the regulation of DNA replication. Single nucleotide polymorphisms in the intron regions of this gene are associated with differential transcriptional activation of the promoter of the neighboring lactase gene and, thereby, influence lactose intolerance in early adulthood. [provided by RefSeq, May 2012]
Function	Acts as component of the MCM2-7 complex (MCM complex) which is the putative replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity. [UniProt]
Calculated Mw	93 kDa
PTM	O-glycosylated (O-GlcNAcylated), in a cell cycle-dependent manner. [UniProt]
Cellular Localization	Nucleus. Note=Binds to chromatin during G1 and detach from it during S phase. [UniProt]

## Images




ARG40479 anti-MCM6 antibody ICC image

Immunocytochemistry: HeLa cells stained with ARG40479 anti-MCM6 antibody.




ARG40479 anti-MCM6 antibody ICC/IF image

Immunofluorescence: A431 cells were blocked with 10% goat serum and then stained with ARG40479 anti-MCM6 antibody (green) at 5 µg/ml dilution, overnight at 4°C. DAPI (blue) for nuclear staining.



ARG40479 anti-MCM6 antibody IHC-Fr image

Immunohistochemistry: Frozen section of Rat intestine tissue. The tissue section was blocked with 10% goat serum. The tissue section was then stained with ARG40479 anti-MCM6 antibody at 1 µg/ml dilution, overnight at 4°C.




ARG40479 anti-MCM6 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human intestinal cancer tissue stained with ARG40479 anti-MCM6 antibody.




ARG40479 anti-MCM6 antibody WB image

Western blot: U87, COLO320, HeLa, MCF7 and Jurkat cell lysates stained with ARG40479 anti-MCM6 antibody.




#### ARG40479 anti-MCM6 antibody WB image

Western blot: 50 µg of sample under reducing conditions. Mouse small intestine and Mouse spleen lysates stained with ARG40479 anti-MCM6 antibody at 0.5 µg/ml dilution, overnight at 4°C.




#### ARG40479 anti-MCM6 antibody FACS image

Flow Cytometry: SiHa cells were blocked with 10% normal goat serum and then stained with ARG40479 anti-MCM6 antibody (blue) at 1 µg/ $10^6$  cells for 30 min at 20°C, followed by incubation with DyLight®488 labelled secondary antibody. Isotype control antibody (green) was rabbit IgG (1 µg/ $10^6$  cells) used under the same conditions. Unlabelled sample (red) was also used as a control.



#### ARG40479 anti-MCM6 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Rat brain stained with ARG40479 anti-MCM6 antibody.



#### ARG40479 anti-MCM6 antibody IHC-Fr image

Immunohistochemistry: Frozen section of Human placenta tissue stained with ARG40479 anti-MCM6 antibody.