

ARG40937 anti-PPP4C antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes PPP4C
Tested Reactivity	Hu, Ms, Rat
Tested Application	IHC-P, IP, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	PPP4C
Species	Human
Immunogen	Recombinant fusion protein corresponding to aa. 1-307 of Human PPP4C (NP_002711.1).
Conjugation	Un-conjugated
Alternate Names	PPP4; PPX; PP-X; Pp4; PPH3; EC 3.1.3.16; Serine/threonine-protein phosphatase 4 catalytic subunit; PP4; PP4C; Protein phosphatase X

Application Instructions

Application table	Application	Dilution
	IHC-P	1:50 - 1:200
	IP	1:50 - 1:200
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Mouse spleen	
Observed Size	36 kDa	

Properties


Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.


Bioinformation

Gene Symbol	PPP4C
Gene Full Name	protein phosphatase 4, catalytic subunit
Function	Protein phosphatase that is involved in many processes such as microtubule organization at centrosomes, maturation of spliceosomal snRNPs, apoptosis, DNA repair, tumor necrosis factor (TNF)-alpha signaling, activation of c-Jun N-terminal kinase MAPK8, regulation of histone acetylation, DNA damage checkpoint signaling, NF-kappa-B activation and cell migration. The PPP4C-PPP4R1 PP4 complex may play a role in dephosphorylation and regulation of HDAC3. The PPP4C-PPP4R2-PPP4R3A PP4 complex specifically dephosphorylates H2AFX phosphorylated on Ser-140 (gamma-H2AFX) generated during DNA replication and required for DNA double strand break repair. Dephosphorylates NDEL1 at CDK1 phosphorylation sites and negatively regulates CDK1 activity in interphase (By similarity). In response to DNA damage, catalyzes RPA2 dephosphorylation, an essential step for DNA repair since it allows the efficient RPA2-mediated recruitment of RAD51 to chromatin. [UniProt]
Calculated Mw	35 kDa
PTM	Methylation at the C-terminal Leu-307 is critical for interactions with regulatory subunits and functions in DNA repair. [UniProt]
Cellular Localization	Cytoplasm. Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. [UniProt]

Images

ARG40937 anti-PPP4C antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human gastric cancer tissue stained with ARG40937 anti-PPP4C antibody at 1:100 dilution.



ARG40937 anti-PPP4C antibody WB image

Western blot: 25 µg of Mouse spleen lysate stained with ARG40937 anti-PPP4C antibody at 1:1000 dilution.

ARG40937 anti-PPP4C antibody IP image

Immunoprecipitation: 200 µg extracts of 293T cells were immunoprecipitated and stained with ARG40937 anti-PPP4C antibody at 1:1000 dilution.

