

Product datasheet

info@arigobio.com

ARG41281 anti-RGS7 antibody

Package: 100 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes RGS7

Tested Reactivity Hu, Ms, Rat

Tested Application WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name RGS7

Species Human

Immunogen Recombinant fusion protein corresponding to aa. 1-300 of Human RGS7 (NP_002915.3).

Conjugation Un-conjugated

Alternate Names RGS7; Regulator of G-protein signaling 7

Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Mouse brain	
Observed Size	58 kDa	

Properties

Form Liquid

Purification Affinity purified.

Buffer PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol RGS7

Gene Full Name regulator of G-protein signaling 7

Function Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving

them into their inactive GDP-bound form. Activity on G(o)-alpha is specifically enhanced by the RGS6/GNG5 dimer. May play a role in synaptic vesicle exocytosis. May play important role in the rapid regulation of neuronal excitability and the cellular responses to short-lived stimulations (By similarity).

[UniProt]

Calculated Mw 58 kDa

PTM Palmitoylated.


Ubiquitinated, leading to rapid proteasomal degradation.

Phosphorylation and subsequent interaction with 14-3-3 proteins inhibits GAP activity. [UniProt]

Cellular Localization Cytoplasm, cytosol. Cytoplasm. Cell membrane. Membrane; Peripheral membrane protein; Cytoplasmic

side. Note=Interaction with PKD1 promotes location at the cell membrane (PubMed:10339594). Interaction with RGS7BP promotes location at the cell membrane (PubMed:15897264). [UniProt]

Images



ARG41281 anti-RGS7 antibody WB image

Western blot: 25 ug of Mouse brain lysate stained with ARG41281 anti-RGS7 antibody at 1:1000 dilution.