

Product datasheet

info@arigobio.com

ARG41306 anti-CSRP3 antibody

Package: 50 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes CSRP3

Tested Reactivity Hu, Rat

Predict Reactivity Ms, Cow, Dog, Gpig, Hrs, Rb

Tested Application WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name CSRP3

Species Human

Immunogen Synthetic peptide around the middle region of Human CSRP3. (within the following region:

QGAGCLSTDTGEHLGLQFQQSPKPARSVTTSNPSKFTAKFGESEKCPRCG)

Conjugation Un-conjugated

Alternate Names CMH12; LIM domain protein, cardiac; Cysteine and glycine-rich protein 3; LMO4; Muscle LIM protein;

MLP; CMD1M; CRP3; Cysteine-rich protein 3; Cardiac LIM protein; CLP

Application Instructions

Predict Reactivity Note Predicted Homology Based On Immunogen Sequence: Cow: 100%; Dog: 100%; Guinea pig: 100%;

Horse: 100%; Mouse: 100%; Rabbit: 100%

Application table Application Dilution

WB 0.2 - 1 μg/ml

Application Note * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations

should be determined by the scientist.

Positive Control Human heart

Properties

Form Liquid

Purification Affinity purified.

Buffer PBS, 0.09% (w/v) Sodium azide and 2% Sucrose.

Preservative 0.09% (w/v) Sodium azide

Stabilizer 2% Sucrose

Concentration Batch dependent: 0.5 - 1 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated

www.arigobio.com arigo, nuts about antibodies 1/2

freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol CSRP3

Gene Full Name cysteine and glycine-rich protein 3 (cardiac LIM protein)

Background This gene encodes a member of the CSRP family of LIM domain proteins, which may be involved in

regulatory processes important for development and cellular differentiation. The LIM/double zincfinger motif found in this protein is found in a group of proteins with critical functions in gene regulation, cell growth, and somatic differentiation. Mutations in this gene are thought to cause heritable forms of hypertrophic cardiomyopathy (HCM) and dilated cardiomyopathy (DCM) in humans. Alternatively spliced transcript variants with different 5' UTR, but encoding the same protein, have

been found for this gene. [provided by RefSeq, Jul 2008]

Function Positive regulator of myogenesis. Plays a crucial and specific role in the organization of cytosolic

structures in cardiomyocytes. Could play a role in mechanical stretch sensing. May be a scaffold protein that promotes the assembly of interacting proteins at Z-line structures. It is essential for calcineurin anchorage to the Z line. Required for stress-induced calcineurin-NFAT activation (By similarity).

[UniProt]

Calculated Mw 21 kDa


PTM Phosphorylated by PKC/PRKCA. [UniProt]

Cellular Localization Nucleus. Cytoplasm. Cytoplasm, cytoskeleton. Cytoplasm, myofibril, sarcomere, Z line. Cytoplasm,

myofibril, sarcomere. Note=Nucleocytoplasmic shuttling protein. Mainly cytoplasmic. In the Z line, found associated with GLRX3 (By similarity). Isoform 2: Cytoplasm, myofibril, sarcomere, Z line.

[UniProt]

Images



ARG41306 anti-CSRP3 antibody WB image

Western blot: Human heart lysate stained with ARG41306 anti-CSRP3 antibody at 0.2-1 $\mu g/ml$ dilution.

Human heart