

ARG43045 anti-Claudin 2 antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes Claudin 2
Tested Reactivity	Hu, Rat
Tested Application	IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Claudin 2
Species	Human
Immunogen	Recombinant protein corresponding to A38-V230 of Human Claudin 2.
Conjugation	Un-conjugated
Alternate Names	Claudin-2; SP82

Application Instructions

Application table	Application	Dilution
	IHC-P	1:200 - 1:1000
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.2% Na ₂ HPO ₄ , 0.9% NaCl, 0.05% Sodium azide and 5% BSA.
Preservative	0.05% Sodium azide
Stabilizer	5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	CLDN2
Gene Full Name	claudin 2
Background	This gene product belongs to the claudin protein family whose members have been identified as major integral membrane proteins localized exclusively at tight junctions. Claudins are expressed in an organ-specific manner and regulate tissue-specific physiologic properties of tight junctions. This protein is expressed in the intestine. Alternatively spliced transcript variants with different 5' untranslated region have been found for this gene. [provided by RefSeq, Jan 2010]
Function	Plays a major role in tight junction-specific obliteration of the intercellular space, through calcium-independent cell-adhesion activity. [UniProt]
Calculated Mw	25 kDa
PTM	The disulfide bond is necessary for pore formation, but is not required for correct protein trafficking. [UniProt]
Cellular Localization	Cell junction, tight junction. Cell membrane; Multi-pass membrane protein. [UniProt]

Images



ARG43045 anti-Claudin 2 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human intestinal cancer tissue stained with ARG43045 anti-Claudin 2 antibody.



ARG43045 anti-Claudin 2 antibody WB image

Western blot: 50 µg of Rat kidney and Rat liver lysates stained with ARG43045 anti-Claudin 2 antibody at 0.5 µg/ml dilution.