

ARG43080 anti-Ku 70 acetyl (Lys539) antibody

Package: 100 µl

Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes Ku 70 acetyl (Lys539)
Tested Reactivity	Hu
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Ku 70
Species	Human
Immunogen	KLH-conjugated synthetic peptide around acetylated Lys539 of Human Ku 70.
Conjugation	Un-conjugated
Alternate Names	DNA repair protein XRCC6; Thyroid-lupus autoantigen; Lupus Ku autoantigen protein p70; EC 4.2.99.-; EC 3.6.4.-; ATP-dependent DNA helicase II 70 kDa subunit; X-ray repair complementing defective repair in Chinese hamster cells 6; CTC box-binding factor 75 kDa subunit; 70 kDa subunit of Ku antigen; CTC75; 5'-deoxyribose-5-phosphate lyase Ku70; KU70; TLAA; 5'-dRP lyase Ku70; CTCBF; ML8; G22P1; X-ray repair cross-complementing protein 6; ATP-dependent DNA helicase 2 subunit 1; Ku70

Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	COLO205 and HEK293T	
Observed Size	70 kDa	

Properties


Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.42% Potassium phosphate (pH 7.3), 0.87% NaCl, 0.01% Sodium azide and 30% Glycerol.
Preservative	0.01% Sodium azide
Stabilizer	30% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	XRCC6
Gene Full Name	X-ray repair complementing defective repair in Chinese hamster cells 6
Background	The p70/p80 autoantigen is a nuclear complex consisting of two subunits with molecular masses of approximately 70 and 80 kDa. The complex functions as a single-stranded DNA-dependent ATP-dependent helicase. The complex may be involved in the repair of nonhomologous DNA ends such as that required for double-strand break repair, transposition, and V(D)J recombination. High levels of autoantibodies to p70 and p80 have been found in some patients with systemic lupus erythematosus. [provided by RefSeq, Jul 2008]
Function	Single-stranded DNA-dependent ATP-dependent helicase. Has a role in chromosome translocation. The DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner. It works in the 3'-5' direction. Binding to DNA may be mediated by XRCC6. Involved in DNA non-homologous end joining (NHEJ) required for double-strand break repair and V(D)J recombination. The XRCC5/6 dimer acts as regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by increasing the affinity of the catalytic subunit PRKDC to DNA by 100-fold. The XRCC5/6 dimer is probably involved in stabilizing broken DNA ends and bringing them together. The assembly of the DNA-PK complex to DNA ends is required for the NHEJ ligation step. Required for osteocalcin gene expression. Probably also acts as a 5'-deoxyribose-5-phosphate lyase (5'-dRP lyase), by catalyzing the beta-elimination of the 5' deoxyribose-5-phosphate at an abasic site near double-strand breaks. 5'-dRP lyase activity allows to 'clean' the termini of abasic sites, a class of nucleotide damage commonly associated with strand breaks, before such broken ends can be joined. The XRCC5/6 dimer together with APEX1 acts as a negative regulator of transcription. Plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP-RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway. [UniProt]
Calculated Mw	70 kDa
PTM	Phosphorylation by PRKDC may enhance helicase activity. Phosphorylation of Ser-51 does not affect DNA repair. [UniProt]
Cellular Localization	Nucleus. Chromosome. [UniProt]

Images

ARG43080 anti-Ku 70 acetyl (Lys539) antibody WB image

Western blot: COLO205 and HEK293T whole cell lysates stained with ARG43080 anti-Ku 70 acetyl (Lys539) antibody.