

ARG44482 anti-INTS14 antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes INTS14
Tested Reactivity	Hu
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	INTS14
Species	Human
Immunogen	Human INTS14 recombinant protein
Conjugation	Un-conjugated
Alternate Names	INTS14; Integrator Complex Subunit 14; C15orf44; VWA9; Von Willebrand Factor A Domain-Containing Protein 9; Von Willebrand Factor A Domain Containing 9; DKFZP564O1664; Chromosome 15 Open Reading Frame 44; UPF0464 Protein C15orf44

Application Instructions

Application table	Application	Dilution
	WB	0.25-0.5 µg/ml
Application Note	The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Affinity purified with Immunogen.
Buffer	0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.05% Sodium azide and 4% Trehalose.
Preservative	0.05% Sodium azide
Stabilizer	4% Trehalose
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	INTS14
Gene Full Name	Integrator Complex Subunit 14
Background	Predicted to be involved in snRNA 3'-end processing. Located in cytoplasm.
Function	Probable component of the Integrator (INT) complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing.
Calculated Mw	57 kDa
PTM	Acetylation
Cellular Localization	Nucleus

Images

ARG44482 anti-INTS14 antibody WB image

Western blot: 293T, Hacat, Jurkat and RT4 stained with ARG44482 anti-INTS14 antibody at 0.5 µg/mL dilution.

