

## ARG44557 anti-CDK1 / CDC2 antibody

Package: 50 µg  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes CDK1 / CDC2
Tested Reactivity	Hu, Ms, Rat
Tested Application	FACS, ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	CDK1 / CDC2
Species	Human
Immunogen	Recombinant Human CDK1 / CDC2 purified from E. coli.
Conjugation	Un-conjugated
Alternate Names	CDK1; Cyclin Dependent Kinase 1; CDC28A; CDC2; Cell Division Cycle 2, G1 To S And G2 To M; Cell Division Control Protein 2 Homolog; Cell Division Protein Kinase 1; Cyclin-Dependent Kinase 1; P34 Protein Kinase; P34CDC2; Cell Cycle Controller CDC2; EC 2.7.11.22; EC 2.7.11.23; CDKN1

### Application Instructions

Application table	Application	Dilution
	FACS	1-3 µg/10 <sup>6</sup> cell
	ICC/IF	2 µg/ml
	IHC-P	0.5-1 µg/ml
	WB	0.1-0.5 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	0.2% Na <sub>2</sub> HPO <sub>4</sub> , 0.9% NaCl, 0.05% Sodium azide and 5% BSA.
Preservative	0.05% Sodium azide
Stabilizer	5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.


#### Note

For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


Gene Symbol	CDK1
Gene Full Name	Cyclin Dependent Kinase 1
Background	The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is a catalytic subunit of the highly conserved protein kinase complex known as M-phase promoting factor (MPF), which is essential for G2/M phase transitions of eukaryotic cell cycle. Mitotic cyclins stably associate with this protein and function as regulatory subunits. The kinase activity of this protein is controlled by cyclin accumulation and destruction through the cell cycle. The phosphorylation and dephosphorylation of this protein also play important regulatory roles in cell cycle control. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2023]
Function	Plays a key role in the control of the eukaryotic cell cycle by modulating the centrosome cycle as well as mitotic onset; promotes G2-M transition via association with multiple interphase cyclins. [UniProt]
Research Area	Cell Biology and Cellular Response antibody; Gene Regulation antibody; Neuroscience antibody
Calculated Mw	34 kDa
PTM	Acetylation, Isopeptide bond, Phosphoprotein, Ubl conjugation. [UniProt]
Cellular Localization	Cytoplasm, Cytoskeleton, Mitochondrion, Nucleus. [UniProt]

## Images



ARG44557 anti-CDK1 / CDC2 antibody IHC-P image

Immunohistochemistry: Human mammary cancer stained with ARG44557 anti-CDK1 / CDC2 antibody at 1 µg/ml dilution.




ARG44557 anti-CDK1 / CDC2 antibody ICC/IF image

Immunofluorescence: U2OS stained with ARG44557 anti-CDK1 / CDC2 antibody at 2 µg/mL dilution.


#### ARG44557 anti-CDK1 / CDC2 antibody WB image

Western blot: HeLa stained with ARG44557 anti-CDK1 / CDC2 antibody at 0.5  $\mu\text{g}/\text{ml}$  dilution.




#### ARG44557 anti-CDK1 / CDC2 antibody FACS image

Flow Cytometry: U937 stained with ARG44557 anti-CDK1 / CDC2 antibody at  $1\mu\text{g}/10^6$  cells dilution.




#### ARG44557 anti-CDK1 / CDC2 antibody IHC-P image


Immunohistochemistry: Rat testis stained with ARG44557 anti-CDK1 / CDC2 antibody at 1  $\mu\text{g}/\text{ml}$  dilution.



#### ARG44557 anti-CDK1 / CDC2 antibody WB image


Western blot: Rat Thymus stained with ARG44557 anti-CDK1 / CDC2 antibody at 0.5  $\mu\text{g}/\text{ml}$  dilution.





ARG44557 anti-CDK1 / CDC2 antibody IHC-P image

Immunohistochemistry: Mouse testis stained with ARG44557 anti-CDK1 / CDC2 antibody at 1 µg/ml dilution.



ARG44557 anti-CDK1 / CDC2 antibody WB image

Western blot: NIH3T3 stained with ARG44557 anti-CDK1 / CDC2 antibody at 0.5 µg/ml dilution.