

ARG44686 anti-ITGB1BP2 / Melusin antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody recognizes ITGB1BP2 / Melusin
Tested Reactivity	Hu, Rat
Tested Application	IP, WB
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Target Name	ITGB1BP2 / Melusin
Species	Human
Conjugation	Un-conjugated
Alternate Names	CHORDC3; Integrin beta-1-binding protein 2; MSTP015; ITGB1BP; Melusin; MELUSIN

Application Instructions

Application table	Application	Dilution
	IP	10 µg/mL
	WB	1 µg/mL
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Protein A purification
Buffer	PBS with 0.09% sodium azide
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	ITGB1BP2
Gene Full Name	integrin beta 1 binding protein (melusin) 2
Background	This gene encodes a protein with two cysteine and histidine-rich (CHORD) domains, PXXP motifs, YXXI/P

motifs, putative SH2 and SH3 domain binding motifs, and an acidic region at the C-terminus that can bind calcium. Two hybrid analysis showed that this protein interacts with the cytoplasmic domain of the beta 1 integrin subunit and is thought to act as a chaperone protein. Studies in the mouse ortholog of this gene indicate that absence of this gene in mouse results in failed cardiac hypertrophy in response to mechanical stress. Alternative splicing results in multiple transcript variants encoding different isoforms, including an isoform that lacks several domains, including one of the CHORD domains. [provided by RefSeq, May 2017]


Function	May play a role during maturation and/or organization of muscles cells. [UniProt]
Calculated Mw	38 kDa
PTM	N-glycosylation enhances cell surface expression and lengthens receptor half-life by preventing degradation in the ER.

Images



ARG44686 anti-ITGB1BP2 / Melusin antibody WB image

Western blot: Rat skeletal stained with ARG44686 anti-ITGB1BP2 / Melusin antibody at 1 µg/mL dilution.



ARG44686 anti-ITGB1BP2 / Melusin antibody IP image

Immunoprecipitation: Rat skeletal lysate immunoprecipitated with 2.5 µg of ARG44686 anti-ITGB1BP2 / Melusin antibody.