

ARG45849 anti-Zebrafish MMP2 antibody

Package: 200 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes MMP2
Tested Reactivity	Zfish
Tested Application	IHC-P
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	MMP2
Species	Zebrafish
Conjugation	Un-conjugated
Alternate Names	MMP2; Matrix Metalloproteinase 2; TBE-1; CLG4A; 72 KDa Type IV Collagenase; Matrix Metalloproteinase-2; EC 3.4.24.24; MMP-2; CLG4; Matrix Metalloproteinase 2 (Gelatinase A, 72kDa Gelatinase, 72kDa Type IV Collagenase); Matrix Metalloproteinase 2 (Gelatinase A, 72kDa Gelatinase, 72kDa Type IV Collagenase); Matrix Metalloproteinase-II; Collagenase Type IV-A; Neutrophil Gelatinase; 72 KDa Gelatinase; Gelatinase A; EC 3.4.24; MMP-II; MONA

Application Instructions


Application table	Application	Dilution
	IHC-P	2-5 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Affinity chromatography purified
Buffer	0.02M PBS, 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% glycerol
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Gene Symbol	MMP2
Gene Full Name	matrix metalloproteinase 2
Background	<p>This gene is a member of the matrix metalloproteinase (MMP) gene family, that are zinc-dependent enzymes capable of cleaving components of the extracellular matrix and molecules involved in signal transduction. The protein encoded by this gene is a gelatinase A, type IV collagenase, that contains three fibronectin type II repeats in its catalytic site that allow binding of denatured type IV and V collagen and elastin. Unlike most MMP family members, activation of this protein can occur on the cell membrane. This enzyme can be activated extracellularly by proteases, or, intracellularly by its S-glutathiolation with no requirement for proteolytical removal of the pro-domain. This protein is thought to be involved in multiple pathways including roles in the nervous system, endometrial menstrual breakdown, regulation of vascularization, and metastasis. Mutations in this gene have been associated with Winchester syndrome and Nodulosis-Arthropathy-Osteolysis (NAO) syndrome. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Oct 2014]</p>
Function	<p>Ubiquitous metalloproteinase that is involved in diverse functions such as remodeling of the vasculature, angiogenesis, tissue repair, tumor invasion, inflammation, and atherosclerotic plaque rupture. As well as degrading extracellular matrix proteins, can also act on several nonmatrix proteins such as big endothelial 1 and beta-type CGRP promoting vasoconstriction. Also cleaves KISS at a Gly- -Leu bond. Appears to have a role in myocardial cell death pathways. Contributes to myocardial oxidative stress by regulating the activity of GSK3beta. Cleaves GSK3beta in vitro. Involved in the formation of the fibrovascular tissues in association with MMP14.</p> <p>PEX, the C-terminal non-catalytic fragment of MMP2, possesses anti-angiogenic and anti-tumor properties and inhibits cell migration and cell adhesion to FGF2 and vitronectin. Ligand for integrinα5β3 on the surface of blood vessels. [UniProt]</p>
Calculated Mw	74 kDa
PTM	Autocatalytic cleavage, Disulfide bond, Glycoprotein, Phosphoprotein, Zymogen. [UniProt]
Cellular Localization	Cytoplasm, Extracellular matrix, Membrane, Mitochondrion, Nucleus, Secreted. [UniProt]

Images



ARG45849 anti-Zebrafish MMP2 antibody IHC-P image

Immunohistochemistry: Zebrafish kidney stained with ARG45849 anti-Zebrafish MMP2 antibody at 2 μ g/ml dilution.