

ARG53783 anti-CD19 antibody [LT19] (PE)

Package: 100 tests Store at: 4°C

Summary

Tested ReactivityHuTested ApplicationFACSSpecificityThe clone LT19 reacts with CD19 (B4), a 95 kDa type I transmembrane glycoprotein (immunoglobulin superfamily) expressed on B lymphocytes and follicular dendritic cells; it is lost on plasma cells.HostMouseClonalityMonoclonalCloneLT19IsotypeIgG1Target NameCD19SpeciesHumanImmunogenDaudi human Burkitt lymphoma cell lineConjugationPE		
Tested ApplicationFACSSpecificityThe clone LT19 reacts with CD19 (B4), a 95 kDa type I transmembrane glycoprotein (immunoglobulin superfamily) expressed on B lymphocytes and follicular dendritic cells; it is lost on plasma cells.HostMouseClonalityMonoclonalCloneLT19IsotypeIgG1Target NameCD19SpeciesHuman Burkitt lymphoma cell lineImmunogenDaudi human Burkitt lymphoma cell lineConjugationPEAlternate NamesDifferentiation antigen CD19; T-cell surface antigen Leu-12; B-lymphocyte antigen CD19; B-lymphocyte	Product Description	PE-conjugated Mouse Monoclonal antibody [LT19] recognizes CD19
SpecificityThe clone LT19 reacts with CD19 (B4), a 95 kDa type I transmembrane glycoprotein (immunoglobulin superfamily) expressed on B lymphocytes and follicular dendritic cells; it is lost on plasma cells.HostMouseClonalityMonoclonalCloneLT19IsotypeIgG1Target NameCD19SpeciesHumanImmunogenDaudi human Burkitt lymphoma cell lineConjugationPEAlternate NamesDifferentiation antigen CD19; T-cell surface antigen Leu-12; B-lymphocyte antigen CD19; B-lymphocyte	Tested Reactivity	Hu
SpeciesHumanImmunogenDaudi human Burkitt lymphoma cell lineConjugationPEAlternate NamesDifferentiation antigen CD19; T-cell surface antigen Leu-12; B-lymphocyte antigen CD19; B-lymphocyte	Tested Application	FACS
ClonalityMonoclonalCloneLT19IsotypeIgG1Target NameCD19SpeciesHumanImmunogenDaudi human Burkitt lymphoma cell lineConjugationPEAlternate NamesDifferentiation antigen CD19; T-cell surface antigen Leu-12; B-lymphocyte antigen CD19; B-lymphocyte	Specificity	
CloneLT19IsotypeIgG1Target NameCD19SpeciesHumanImmunogenDaudi human Burkitt lymphoma cell lineConjugationPEAlternate NamesDifferentiation antigen CD19; T-cell surface antigen Leu-12; B-lymphocyte antigen CD19; B-lymphocyte	Host	Mouse
IsotypeIgG1Target NameCD19SpeciesHumanImmunogenDaudi human Burkitt lymphoma cell lineConjugationPEAlternate NamesDifferentiation antigen CD19; T-cell surface antigen Leu-12; B-lymphocyte antigen CD19; B-lymphocyte	Clonality	Monoclonal
Target NameCD19SpeciesHumanImmunogenDaudi human Burkitt lymphoma cell lineConjugationPEAlternate NamesDifferentiation antigen CD19; T-cell surface antigen Leu-12; B-lymphocyte antigen CD19; B-lymphocyte	Clone	LT19
Species Human Immunogen Daudi human Burkitt lymphoma cell line Conjugation PE Alternate Names Differentiation antigen CD19; T-cell surface antigen Leu-12; B-lymphocyte antigen CD19; B-lymphocyte	Isotype	lgG1
Immunogen Daudi human Burkitt lymphoma cell line Conjugation PE Alternate Names Differentiation antigen CD19; T-cell surface antigen Leu-12; B-lymphocyte antigen CD19; B-lymphocyte	Target Name	CD19
Conjugation PE Alternate Names Differentiation antigen CD19; T-cell surface antigen Leu-12; B-lymphocyte antigen CD19; B-lymphocyte	Species	Human
Alternate Names Differentiation antigen CD19; T-cell surface antigen Leu-12; B-lymphocyte antigen CD19; B-lymphocyte	Immunogen	Daudi human Burkitt lymphoma cell line
	Conjugation	PE
	Alternate Names	

Application Instructions

Application table	Application	Dilution
	FACS	20 μl / 10^6 cells
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Liquid
The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
PBS, 15 mM Sodium azide and 0.2% (w/v) high-grade protease free BSA
15 mM Sodium azide
0.2% (w/v) high-grade protease free BSA
Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Database links	GenelD: 930 Human
	Swiss-port # P15391 Human
Gene Symbol	CD19
Gene Full Name	CD19 molecule
Background	CD19: Lymphocytes proliferate and differentiate in response to various concentrations of different antigens. The ability of the B cell to respond in a specific, yet sensitive manner to the various antigens is achieved with the use of low-affinity antigen receptors. This gene encodes a cell surface molecule which assembles with the antigen receptor of B lymphocytes in order to decrease the threshold for antigen receptor-dependent stimulation. [provided by RefSeq, Jul 2008]
Function	CD19 functions as coreceptor for the B-cell antigen receptor complex (BCR) on B-lymphocytes. Decreases the threshold for activation of downstream signaling pathways and for triggering B-cell responses to antigens (PubMed:2463100, PubMed:1373518, PubMed:16672701). Activates signaling pathways that lead to the activation of phosphatidylinositol 3-kinase and the mobilization of intracellular Ca(2+) stores (PubMed:9382888, PubMed:9317126, PubMed:12387743, PubMed:16672701). Is not required for early steps during B cell differentiation in the blood marrow (PubMed:9317126). Required for normal differentiation of B-1 cells. Required for normal B cell differentiation and proliferation in response to antigen challenges (PubMed:2463100, PubMed:1373518). Required for normal levels of serum immunoglobulins, and for production of high- affinity antibodies in response to antigen challenge (PubMed:9317126, PubMed:12387743, PubMed:16672701). [UniProt]
Highlight	Related products: <u>CD19 antibodies;</u> <u>CD19 ELISA Kits;</u> <u>CD19 Duos / Panels;</u> <u>Anti-Mouse IgG secondary antibodies;</u> Related news: <u>Tumor-Infiltrating Lymphocytes (TILs)</u>
Research Area	Developmental Biology antibody; Immune System antibody; Lymphocyte Marker antibody; B cell Marker antibody; Pro-B Cell Marker antibody; Pre-B Cell Marker antibody; Immature B Cell Marker antibody; Follicular dendritic cells antibody
Calculated Mw	61 kDa
РТМ	Phosphorylated on serine and threonine upon DNA damage, probably by ATM or ATR. Phosphorylated on tyrosine following B-cell activation. Phosphorylated on tyrosine residues by LYN.

Bioinformation

Images



ARG53783 anti-CD19 antibody [LT19] (PE) FACS image

Flow Cytometry: Human peripheral blood cells stained with ARG53783 anti-CD19 antibody [LT19] (PE).