

ARG53784 anti-CD19 antibody [LT19] (PerCP)

Package: 100 tests Store at: 4°C

Summary

Product Description	PerCP-conjugated Mouse Monoclonal antibody [LT19] recognizes CD19
Tested Reactivity	Hu
Tested Application	FACS
Specificity	The clone LT19 reacts with CD19 (B4), a 95 kDa type I transmembrane glycoprotein (immunoglobulin superfamily) expressed on B lymphocytes and follicular dendritic cells; it is lost on plasma cells.
Host	Mouse
Clonality	Monoclonal
Clone	LT19
Isotype	lgG1
Target Name	CD19
Species	Human
Immunogen	Daudi human Burkitt lymphoma cell line
Conjugation	PerCP
Alternate Names	Differentiation antigen CD19; T-cell surface antigen Leu-12; B-lymphocyte antigen CD19; B-lymphocyte surface antigen B4; B4; CD antigen CD19; CVID3

Application Instructions

Application table	Application	Dilution
	FACS	10 μl / 10^6 cells
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. N reconstitution is necessary.BufferPBS, 15 mM Sodium azide and 0.2% (w/v) high-grade protease free BSAPreservative15 mM Sodium azideStabilizer0.2% (w/v) high-grade protease free BSAStorage instructionAliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be	Form	Liquid
Preservative 15 mM Sodium azide Stabilizer 0.2% (w/v) high-grade protease free BSA Storage instruction Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be	Purification Note	The purified antibody is conjugated with Peridinin-chlorophyll-protein complex (PerCP) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.
Stabilizer0.2% (w/v) high-grade protease free BSAStorage instructionAliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be	Buffer	PBS, 15 mM Sodium azide and 0.2% (w/v) high-grade protease free BSA
Storage instructionAliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be	Preservative	15 mM Sodium azide
repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be	Stabilizer	0.2% (w/v) high-grade protease free BSA
	Storage instruction	repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be

Database links	GeneID: 930 Human
	Swiss-port # P15391 Human
Gene Symbol	CD19
Gene Full Name	CD19 molecule
Background	CD19: Lymphocytes proliferate and differentiate in response to various concentrations of different antigens. The ability of the B cell to respond in a specific, yet sensitive manner to the various antigens is achieved with the use of low-affinity antigen receptors. This gene encodes a cell surface molecule which assembles with the antigen receptor of B lymphocytes in order to decrease the threshold for antigen receptor-dependent stimulation. [provided by RefSeq, Jul 2008]
Function	CD19 functions as coreceptor for the B-cell antigen receptor complex (BCR) on B-lymphocytes. Decreases the threshold for activation of downstream signaling pathways and for triggering B-cell responses to antigens (PubMed:2463100, PubMed:1373518, PubMed:16672701). Activates signaling pathways that lead to the activation of phosphatidylinositol 3-kinase and the mobilization of intracellular Ca(2+) stores (PubMed:9382888, PubMed:9317126, PubMed:12387743, PubMed:16672701). Is not required for early steps during B cell differentiation in the blood marrow (PubMed:9317126). Required for normal differentiation of B-1 cells. Required for normal B cell differentiation and proliferation in response to antigen challenges (PubMed:2463100, PubMed:1373518). Required for normal levels of serum immunoglobulins, and for production of high- affinity antibodies in response to antigen challenge (PubMed:9317126, PubMed:12387743, PubMed:16672701). [UniProt]
Highlight	Related products: <u>CD19 antibodies; CD19 ELISA Kits; CD19 Duos / Panels; Anti-Mouse IgG secondary antibodies;</u> Related news: <u>Tumor-Infiltrating Lymphocytes (TILs)</u>
Research Area	Developmental Biology antibody; Immune System antibody; Lymphocyte Marker antibody; B cell Marker antibody; Pro-B Cell Marker antibody; Pre-B Cell Marker antibody; Immature B Cell Marker antibody; Follicular dendritic cells antibody
Calculated Mw	61 kDa
PTM	Phosphorylated on serine and threonine upon DNA damage, probably by ATM or ATR. Phosphorylated on tyrosine following B-cell activation. Phosphorylated on tyrosine residues by LYN.

Bioinformation

Images



ARG53784 anti-CD19 antibody [LT19] (PerCP) FACS image

Flow Cytometry: Human peripheral blood cells stained with ARG53784 anti-CD19 antibody [LT19] (PerCP).