

Product datasheet

info@arigobio.com

ARG55063 anti-Calpain 2 antibody [1381CT669.7.66.71]

Package: 100 μl Store at: -20°C

Summary

Product Description Mouse Monoclonal antibody recognizes Calpain 2

Tested Reactivity Hu, Ms

Tested Application FACS, IHC-P, WB

Host Mouse

Clonality Monoclonal

Clone 1381CT669.7.66.71

Isotype IgG2b

Target Name Calpain 2

Species Human

Immunogen KLH-conjugated synthetic peptide of Human Calpain 2.

Conjugation Un-conjugated

Alternate Names CANP2; Calpain-2 large subunit; Calpain-2 catalytic subunit; EC 3.4.22.53; Millimolar-calpain; M-calpain;

mCANP; Calpain large polypeptide L2; Calpain M-type; CANP 2; CANPL2; CANPMI; Calcium-activated

neutral proteinase 2

Application Instructions

Application table	Application	Dilution
	FACS	1:100
	IHC-P	Assay-dependent
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	A549	

Properties

Form Liquid

Purification Purification with Protein G.

Buffer PBS and 0.09% (W/V) Sodium azide

Preservative 0.09% (W/V) Sodium azide

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Bioinformation

Database links GeneID: 12334 Mouse

GeneID: 824 Human

Swiss-port # O08529 Mouse

Swiss-port # P17655 Human

Gene Symbol CAPN2

Gene Full Name calpain 2, (m/II) large subunit

Background The calpains, calcium-activated neutral proteases, are nonlysosomal, intracellular cysteine proteases.

The mammalian calpains include ubiquitous, stomach-specific, and muscle-specific proteins. The ubiquitous enzymes consist of heterodimers with distinct large, catalytic subunits associated with a common small, regulatory subunit. This gene encodes the large subunit of the ubiquitous enzyme, calpain 2. Multiple heterogeneous transcriptional start sites in the 5' UTR have been reported. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar

2009]

Function Calcium-regulated non-lysosomal thiol-protease which catalyze limited proteolysis of substrates

involved in cytoskeletal remodeling and signal transduction. Proteolytically cleaves MYOC at 'Arg-226'

(PubMed:17650508). [UniProt]


Research Area Cancer antibody; Cell Biology and Cellular Response antibody; Metabolism antibody; Signaling

Transduction antibody

Calculated Mw 80 kDa


Cellular Localization Cytoplasm. Cell membrane. Note=Translocates to the plasma membrane upon Ca(2+) binding

Images




ARG55063 anti-Calpain 2 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human stomach tissue stained with ARG55063 anti-Calpain 2 antibody at 1:25 dilution.



ARG55063 anti-Calpain 2 antibody WB image

Western blot: 35 μg of A549 cell lysate stained with ARG55063 anti-Calpain 2 antibody at 1:1000 dilution.



ARG55063 anti-Calpain 2 antibody FACS image

Flow Cytometry: U-87 MG cells stained with ARG55063 anti-Calpain 2 antibody (right histogram) at 1:100 dilution or isotype control antibody (left histogram), followed by incubation with Alexa Fluor® 488 labelled secondary antibody.