

ARG55128 anti-ADA antibody [608CT2.1.3]

Package: 100 μl Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody recognizes ADA
Tested Reactivity	Hu
Tested Application	WB
Host	Mouse
Clonality	Monoclonal
Clone	608CT2.1.3
Isotype	lgG1
Target Name	ADA
Species	Human
Immunogen	KLH-conjugated synthetic peptide corresponding to aa. 287-314 (C-terminus) of Human ADA.
Conjugation	Un-conjugated
Alternate Names	Adenosine deaminase; EC 3.5.4.4; Adenosine aminohydrolase

Application Instructions

Application table	Application	Dilution
	WB	1:100 - 1:250
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Jurkat	

Properties

Form	Liquid
Purification	Purification with Protein G.
Buffer	PBS and 0.09% (W/V) Sodium azide
Preservative	0.09% (W/V) Sodium azide
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GenelD: 100 Human
	Swiss-port # P00813 Human
Gene Symbol	ADA
Gene Full Name	adenosine deaminase
Background	This gene encodes an enzyme that catalyzes the hydrolysis of adenosine to inosine. Various mutations have been described for this gene and have been linked to human diseases. Deficiency in this enzyme causes a form of severe combined immunodeficiency disease (SCID), in which there is dysfunction of both B and T lymphocytes with impaired cellular immunity and decreased production of immunoglobulins, whereas elevated levels of this enzyme have been associated with congenital hemolytic anemia.
Function	Catalyzes the hydrolytic deamination of adenosine and 2-deoxyadenosine. Plays an important role in purine metabolism and in adenosine homeostasis. Modulates signaling by extracellular adenosine, and so contributes indirectly to cellular signaling events. Acts as a positive regulator of T-cell coactivation, by binding DPP4. Its interaction with DPP4 regulates lymphocyte-epithelial cell adhesion. [UniProt]
Research Area	Cancer antibody; Gene Regulation antibody; Metabolism antibody
Calculated Mw	41 kDa

Images

