

# Product datasheet

info@arigobio.com

ARG55765 anti-RPS21 antibody

Package: 100 μl Store at: -20°C

## **Summary**

Product Description Rabbit Polyclonal antibody recognizes RPS21

Tested Reactivity Hu

Predict Reactivity Ms, Rat, Bov, Pig

Tested Application WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name RPS21

Species Human

Immunogen KLH-conjugated synthetic peptide corresponding to aa. 44-71 (C-terminus) of Human RPS21.

Conjugation Un-conjugated

Alternate Names HLDF; S21; 40S ribosomal protein S21

## **Application Instructions**

Application table	Application	Dilution
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	NCI-H460	

### **Properties**

Form Liquid

**Purification** Purification with Protein A and immunogen peptide.

Buffer PBS and 0.09% (W/V) Sodium azide.

Preservative 0.09% (W/V) Sodium azide.

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Database links GeneID: 6227 Human

Swiss-port # P63220 Human

Gene Symbol RPS21

Gene Full Name ribosomal protein S21

Background Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S

subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 40S subunit. The protein belongs to the S21E family of ribosomal proteins. It is located in the cytoplasm. Alternative splice variants that encode different protein isoforms have been described, but their existence has not been verified. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. [provided by RefSeq, Jul 2008]

Calculated Mw 9 kDa

## **Images**

