

## ARG55992 anti-basic Cytokeratin antibody [AE3]

Package: 50 µg  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody [AE3] recognizes basic Cytokeratin
Tested Reactivity	Hu, Ms, Rat
Tested Application	FACS, ICC/IF, IHC-P, WB
Host	Mouse
Clonality	Monoclonal
Clone	AE3
Isotype	IgG1
Target Name	basic Cytokeratin
Species	Human
Immunogen	Solubilized cytokeratin extract from Human stratum corneum.
Conjugation	Un-conjugated
Alternate Names	67 kDa cytokeratin; KRT1A; EPPK; Keratin, type II cytoskeletal 1; EHK1; K1; Keratin-1; CK1; Cytokeratin-1; Hair alpha protein; Type-II keratin Kb1; CK-1; NEPPK; EHK

### Application Instructions

Application table	Application	Dilution
	FACS	0.5 - 1 µg/10 <sup>6</sup> cells
	ICC/IF	1 - 5 µg/ml
	IHC-P	1 - 5 µg/ml
	WB	0.5 - 1 µg/ml
Application Note	IHC-P: Antigen Retrieval: boil tissue sections in 10 mM Citrate buffer (pH 6.0) for 10-20 min, followed by cooling at RT for 20 min. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties


Form	Liquid
Purification	Purification with Protein G.
Buffer	PBS (pH 7.4), 0.05% Sodium azide and 0.1 mg/ml BSA
Preservative	0.05% Sodium azide
Stabilizer	0.1 mg/ml BSA
Concentration	0.2 mg/ml

<b>Storage instruction</b>	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
<b>Note</b>	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

<b>Gene Symbol</b>	KRT1
<b>Gene Full Name</b>	keratin 1, type II
<b>Background</b>	The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins which are arranged in pairs of heterotypic keratin chains coexpressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the spinous and granular layers of the epidermis with family member KRT10 and mutations in these genes have been associated with bullous congenital ichthyosiform erythroderma. The type II cytokeratins are clustered in a region of chromosome 12q12-q13. [provided by RefSeq, Jul 2008]
<b>Function</b>	May regulate the activity of kinases such as PKC and SRC via binding to integrin beta-1 (ITB1) and the receptor of activated protein kinase C (RACK1/GNB2L1). In complex with C1QBP is a high affinity receptor for kininogen-1/HMWK. [UniProt]
<b>Research Area</b>	Signaling Transduction antibody; Carcinoma Marker antibody
<b>Calculated Mw</b>	66 kDa
<b>PTM</b>	Undergoes deimination of some arginine residues (citrullination).
<b>Cellular Localization</b>	Cytoplasmic

## Images



ARG55992 anti-basic Cytokeratin antibody [AE3] IHC-P image

Immunohistochemistry: Formalin-fixed, paraffin-embedded skin stained with ARG55992 anti-basic Cytokeratin antibody [AE3].