

ARG56254 anti-CTCF antibody

Package: 100 μl Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes CTCF
Tested Reactivity	Hu, Ms, Rat
Tested Application	ChIP, ICC/IF, IHC-P, IP, WB
Host	Rabbit
Clonality	Polyclonal
lsotype	lgG
Target Name	CTCF
Species	Human
Immunogen	Recombinant protein of Human CTCF
Conjugation	Un-conjugated
Alternate Names	CCCTC-binding factor; 11-zinc finger protein; Transcriptional repressor CTCF; MRD21; CTCFL paralog

Application Instructions

Application table	Application	Dilution
	ChIP	Assay-dependent
	ICC/IF	1:50 - 1:200
	IHC-P	1:50 - 1:200
	IP	Assay-dependent
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	U251	

Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Bioinformation

Gene Symbol	CTCF
Gene Full Name	CCCTC-binding factor (zinc finger protein)
Background	This gene is a member of the BORIS + CTCF gene family and encodes a transcriptional regulator protein with 11 highly conserved zinc finger (ZF) domains. This nuclear protein is able to use different combinations of the ZF domains to bind different DNA target sequences and proteins. Depending upon the context of the site, the protein can bind a histone acetyltransferase (HAT)-containing complex and function as a transcriptional activator or bind a histone deacetylase (HDAC)-containing complex and function as a transcriptional repressor. If the protein is bound to a transcriptional insulator element, it can block communication between enhancers and upstream promoters, thereby regulating imprinted expression. Mutations in this gene have been associated with invasive breast cancers, prostate cancers, and Wilms' tumors. Alternatively spliced transcript variants encoding different isoforms have been found
	for this gene. [provided by RefSeq, Jul 2010]
Function	Chromatin binding factor that binds to DNA sequence specific sites. Involved in transcriptional regulation by binding to chromatin insulators and preventing interaction between promoter and nearby enhancers and silencers. Acts as transcriptional repressor binding to promoters of vertebrate MYC gene and BAG1 gene. Also binds to the PLK and PIM1 promoters. Acts as a transcriptional activator of APP. Regulates APOA1/C3/A4/A5 gene cluster and controls MHC class II gene expression. Plays an essential role in oocyte and preimplantation embryo development by activating or repressing transcription. Seems to act as tumor suppressor. Plays a critical role in the epigenetic regulation. Participates in the allele-specific gene expression at the imprinted IGF2/H19 gene locus. On the maternal allele, binding within the H19 imprinting control region (ICR) mediates maternally inherited higher-order chromatin conformation to restrict enhancer access to IGF2. Plays a critical role in gene silencing over considerable distances in the genome. Preferentially interacts with unmethylated DNA, preventing spreading of CpG methylation. Plays a important role in chromatin remodeling. Can dimerize when it is bound to different DNA sequences, mediating long-range chromatin looping. Mediates interchromosomal association between IGF2/H19 and WSB1/NF1 and may direct distant DNA segments to a common transcription factory. Causes local loss of histone acetylation and gain of histone methylation in the beta-globin locus, without affecting transcription. When bound to chromatin, it provides an anchor point for nucleosomes positioning. Seems to be essential for homologous X-chromosome pairing. May participate with Tsix in establishing a regulatable epigenetic switch for X chromosome inactivation. May play a role in preventing the propagation of stable methylation at the escape genes from X- inactivation. Involved in sister chromatid cohesion. Associates with both centromeres and chromosomal arms during metaphase and
Coloridate d Mari	required for cohesin localization to CTCF sites. Regulates asynchronous replication of IGF2/H19. [UniProt]
	83 KUB Sumpulated on Lys 74 and Lys 690, sumpulation of CTCE contributes to the represence function of CTCE on
PTM	Sumoviated on Lys-74 and Lys-689; sumoviation of CTCF contributes to the repressive function of CTCF on the MYC P2 promoter.

Images



ARG56254 anti-CTCF antibody IHC-P image

Immunohistochemistry: Human colon carcinoma stained with ARG56254 anti-CTCF antibody at 1:50 dilution.





ARG56254 anti-CTCF antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human lung cancer stained with ARG56254 anti-CTCF antibody at 1:100 dilution.

ARG56254 anti-CTCF antibody WB image

Western blot: U251 cell lysate stained with ARG56254 anti-CTCF antibody.