

ARG56467 anti-H-FABP / Cardiac FABP antibody [CC68]

Package: 100 µg
Store at: -20°C

Summary

Product Description	Mouse Monoclonal antibody [CC68] recognizes H-FABP / Cardiac FABP
Tested Reactivity	Hu, Rat
Species Does Not React With	Ms
Tested Application	WB
Specificity	This antibody does not react to Mouse FABP1, FABP2, FABP3, FABP4 and FABP5.
Host	Mouse
Clonality	Monoclonal
Clone	CC68
Isotype	IgG2b
Target Name	H-FABP / Cardiac FABP
Species	Human
Immunogen	Synthetic peptide around aa. 44-55 of Human Cardiac FABP.
Conjugation	Un-conjugated
Alternate Names	FABP11; H-FABP; O-FABP; Heart-type fatty acid-binding protein; MDGI; Fatty acid-binding protein 3; Muscle fatty acid-binding protein; Mammary-derived growth inhibitor; Fatty acid-binding protein, heart; M-FABP

Application Instructions

Application table	Application	Dilution
	WB	1:200
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Purified
Buffer	PBS (pH 7.2), 0.02% Sodium azide, 50% Glycerol and 0.1% BSA.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol and 0.1% BSA
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 2170 Human GeneID: 79131 Rat Swiss-port # P05413 Human Swiss-port # P07483 Rat
Gene Symbol	FABP3
Gene Full Name	fatty acid binding protein 3, muscle and heart
Background	The intracellular fatty acid-binding proteins (FABPs) belongs to a multigene family. FABPs are divided into at least three distinct types, namely the hepatic-, intestinal- and cardiac-type. They form 14-15 kDa proteins and are thought to participate in the uptake, intracellular metabolism and/or transport of long-chain fatty acids. They may also be responsible in the modulation of cell growth and proliferation. Fatty acid-binding protein 3 gene contains four exons and its function is to arrest growth of mammary epithelial cells. This gene is a candidate tumor suppressor gene for human breast cancer. [provided by RefSeq, Jul 2008]
Function	FABP are thought to play a role in the intracellular transport of long-chain fatty acids and their acyl-CoA esters. [UniProt]
Calculated Mw	15 kDa