

ARG56754 anti-IL1 beta antibody (Biotin)

Package: 50 µg
Store at: 4°C

Summary

Product Description	Biotin-conjugated Rabbit Polyclonal antibody recognizes IL1 beta
Tested Reactivity	Ms
Tested Application	ELISA, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	IL1 beta
Species	Mouse
Immunogen	E.coli derived Recombinant Mouse IL-1β. (MVPIRQLHYR LRDEQQKSLV LSDPYELKAL HLNQINQQ VIFSMSFVQG EPSNDKIPVA LGLKGKNLYL SCVMKDGTP T LQLESVDPKQ YPKKKMEKRF VFNKIEVKSK VEFESAEPN WYISTSQAEH KPVFLGNNSG QDIIDFTMES VSS)
Conjugation	Biotin
Alternate Names	Interleukin-1 beta; IL1-BETA; IL-1; IL-1 beta; Catabolin; IL1F2

Application Instructions


Application table	Application	Dilution
	ELISA	Direct: 0.25 - 1.0 µg/ml Sandwich: 0.25 - 1.0 µg/ml with ARG56644 as a capture antibody
	WB	0.1 - 0.2 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Purified by affinity chromatography.
Buffer	PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


Database links	GeneID: 16176 Mouse Swiss-port # P10749 Mouse
Gene Symbol	Il1b
Gene Full Name	interleukin 1 beta
Background	IL1 beta protein is a member of the interleukin 1 cytokine family. This cytokine is produced by activated macrophages as a proprotein, which is proteolytically processed to its active form by caspase 1 (CASP1/ICE). This cytokine is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. The induction of cyclooxygenase-2 (PTGS2/COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. [provided by RefSeq, Jul 2008]
Function	IL1 beta is a potent proinflammatory cytokine. Initially discovered as the major endogenous pyrogen, induces prostaglandin synthesis, neutrophil influx and activation, T-cell activation and cytokine production, B-cell activation and antibody production, and fibroblast proliferation and collagen production. Promotes Th17 differentiation of T-cells. Synergizes with IL12/interleukin-12 to induce IFNG synthesis from T-helper 1 (Th1) cells (PubMed:10653850). [UniProt]
Highlight	Related products: IL1 beta antibodies ; IL1 beta ELISA Kits ; IL1 beta Duos / Panels ; IL1 beta recombinant proteins ; Anti-Rabbit IgG secondary antibodies ; Related news: HMGB1 in inflammation Inflammatory Cytokines Exploring Antiviral Immune Response RIP1 activation and pathogenesis of NASH
Research Area	Pyroptosis Study antibody
Calculated Mw	31 kDa
PTM	Activation of the IL1B precursor involves a CASP1-catalyzed proteolytic cleavage. Processing and secretion are temporarily associated.

Images



ARG56754 anti-IL1 beta antibody (Biotin) standard curve image

Direct ELISA: ARG56754 anti-IL1 beta antibody (Biotin) at 0.25 - 1.0 µg/ml results of a typical standard run with optical density.



ARG56754 anti-IL1 beta antibody (Biotin) standard curve image

Sandwich ELISA: ARG56754 anti-IL1 beta antibody (Biotin) as a detection antibody at 0.25 - 1.0 $\mu\text{g/ml}$ combined with ARG56644 anti-IL1 beta antibody as a capture antibody. Results of a typical standard run with optical density.