

Product datasheet

info@arigobio.com

ARG57003 anti-Cytokeratin 23 antibody [2F6]

Package: 50 μl Store at: -20°C

Summary

Product Description Mouse Monoclonal antibody [2F6] recognizes Cytokeratin 23

Tested Reactivity Hu

Tested Application FACS, ICC/IF, WB

Host Mouse

Clonality Monoclonal

2F6 Clone

Isotype IgG1, kappa **Target Name** Cytokeratin 23

Species Human

Immunogen Recombinant fragment corresponding to aa. 271-422 of Human Cytokeratin 23.

Conjugation Un-conjugated

K23; CK23; CK-23; Keratin-23; HAIK1; Cytokeratin-23; Keratin, type I cytoskeletal 23 **Alternate Names**

Application Instructions

Application table	Application	Dilution
	FACS	2 - 5 μg/10^6 cells
	ICC/IF	1:100
	WB	1:500 - 1:3000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form Liquid

Purification Purification with Protein G.

Buffer PBS (pH 7.4), 0.02% Sodium azide and 10% Glycerol.

1 mg/ml

Preservative 0.02% Sodium azide

Stabilizer 10% Glycerol Concentration

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Bioinformation

Database links <u>GeneID: 25984 Human</u>

Swiss-port # Q9C075 Human

Gene Symbol KRT23

Gene Full Name keratin 23, type I


Background The protein encoded by this gene is a member of the keratin family. The keratins are intermediate

filament proteins responsible for the structural integrity of epithelial cells and are subdivided into cytokeratins and hair keratins. The type I cytokeratins consist of acidic proteins which are arranged in pairs of heterotypic keratin chains. The type I cytokeratin genes are clustered in a region of chromosome 17q12-q21. Alternative splicing results in multiple transcript variants. [provided by

RefSeq, Sep 2013]


Calculated Mw 48 kDa

Images




ARG57003 anti-Cytokeratin 23 antibody [2F6] ICC/IF image

Immunoflorescense: Hep3B cells stained with ARG57003 anti-Cytokeratin 23 antibody [2F6] at 1:100 (green). DAPI (blue) for nucleus staining.



ARG57003 anti-Cytokeratin 23 antibody [2F6] FACS image

Flow Cytometry: Hep3B cell line stained with ARG57003 anti-Cytokeratin 23 antibody [2F6] at 2-5 μ g for 1x10^6 cells. Secondary antibody: Goat anti-Mouse IgG Alexa fluor 488 conjugate.



ARG57003 anti-Cytokeratin 23 antibody [2F6] WB image

Western blot: 40 μg of HeLa and HepG2 cell lysates stained with ARG57003 anti-Cytokeratin 23 antibody [2F6] at 1:3000 dilution.