

Product datasheet

info@arigobio.com

ARG57503 anti-HSPA13 / STCH antibody [2F6]

Package: 50 μl Store at: -20°C

Summary

Product Description Mouse Monoclonal antibody [2F6] recognizes HSPA13 / STCH

Tested Reactivity Hu

Tested Application ICC/IF, WB
Host Mouse

Clonality Monoclonal

Clone 2F6

Isotype IgG1, kappa

Target Name HSPA13 / STCH

Species Human

Immunogen Recombinant Human HSPA13 (aa. 23-471) purified from E. coli.

Conjugation Un-conjugated

Alternate Names STCH; Heat shock 70 kDa protein 13; Microsomal stress-70 protein ATPase core; Stress-70 protein

chaperone microsome-associated 60 kDa protein

Application Instructions

| Application table | Application | Dilution |
|-------------------|--|----------|
| | ICC/IF | 1:100 |
| | WB | 1:1000 |
| Application Note | * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. | |

Properties

Form Liquid

Purification Purification with Protein A.

Buffer PBS (pH 7.4), 0.02% Sodium azide and 10% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 10% Glycerol

Concentration 1 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol HSPA13

Gene Full Name heat shock protein 70kDa family, member 13

Background The protein encoded by this gene is a member of the heat shock protein 70 family and is found


associated with microsomes. Members of this protein family play a role in the processing of cytosolic and secretory proteins, as well as in the removal of denatured or incorrectly-folded proteins. The encoded protein contains an ATPase domain and has been shown to associate with a ubiquitin-like

protein. [provided by RefSeq, Jul 2008]

Function Has peptide-independent ATPase activity. [UniProt]


Calculated Mw 52 kDa

Images



ARG57503 anti-HSPA13 / STCH antibody [2F6] ICC/IF image

Immunoflorescense: Hep3B cells stained with ARG57503 anti-HSPA13 / STCH antibody [2F6] at 1:100 dilution. The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).



ARG57503 anti-HSPA13 / STCH antibody [2F6] WB image

Western blot: 40 μg of 293T, K562 and MCF7 cell lysates stained with ARG57503 anti-HSPA13 / STCH antibody [2F6] at 1:1000 dilution.