

Product datasheet

info@arigobio.com

ARG58990 anti-CHRAC1 antibody

Package: 50 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes CHRAC1

Tested Reactivity Hu, Ms, Rat

Tested Application WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name CHRAC1

Species Human

Immunogen Recombinant fusion protein corresponding to aa. 1-131 of Human CHRAC1 (NP_059140.1).

Conjugation Un-conjugated

Alternate Names CHRAC-1; Chromatin accessibility complex protein 1; CHRAC-15; YCL1; CHARC1; DNA polymerase

epsilon subunit p15; CHARC15; HuCHRAC15; CHRAC15; Chromatin accessibility complex 15 kDa protein

Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HeLa	
Observed Size	~ 16 kDa	

Properties

Form Liquid

Purification Affinity purified.

Buffer PBS (pH 7.3), 0.02% Sodium azide and 50% Glycerol.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol CHRAC1

Gene Full Name chromatin accessibility complex 1

Background CHRAC1 is a histone-fold protein that interacts with other histone-fold proteins to bind DNA in a

sequence-independent manner. These histone-fold protein dimers combine within larger enzymatic

complexes for DNA transcription, replication, and packaging.[supplied by OMIM, Apr 2004]

Function Forms a complex with DNA polymerase epsilon subunit POLE3 and binds naked DNA, which is then


incorporated into chromatin, aided by the nucleosome remodeling activity of ISWI/SNF2H and ACF1.

[UniProt]

Calculated Mw 15 kDa

Cellular Localization Nucleus. [UniProt]

Images



ARG58990 anti-CHRAC1 antibody WB image

Western blot: 25 μg of HeLa cell lysate stained with ARG58990 anti-CHRAC1 antibody at 1:1000 dilution.