

Product datasheet

info@arigobio.com

ARG59533 anti-Kisspeptin antibody

Package: 100 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes Kisspeptin

Tested Reactivity Hu

Tested Application IHC-P, WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name Kisspeptin
Species Human

Immunogen KLH-conjugated synthetic peptide between aa. 20-52 of Human Kisspeptin.

Conjugation Un-conjugated

Alternate Names Kisspeptin-1; Kisspeptin-54; HH13; KiSS-1; Metastasis-suppressor KiSS-1

Application Instructions

Application table	Application	Dilution
	IHC-P	1:25
	WB	1:1000 - 1:2000
Application Note	IHC-P: Antigen Retrieval: Heat mediation was performed in Citrate buffer (pH 6.0). * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Human skeletal muscle	

Properties

Form Liquid

Purification Purification with Protein A and immunogen peptide.

Buffer PBS and 0.09% (W/V) Sodium azide.

Preservative 0.09% (W/V) Sodium azide.

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol KISS1

Gene Full Name KiSS-1 metastasis-suppressor

Background This gene is a metastasis suppressor gene that suppresses metastases of melanomas and breast

carcinomas without affecting tumorigenicity. The encoded protein may inhibit chemotaxis and invasion and thereby attenuate metastasis in malignant melanomas. Studies suggest a putative role in the regulation of events downstream of cell-matrix adhesion, perhaps involving cytoskeletal reorganization. A protein product of this gene, kisspeptin, stimulates gonadotropin-releasing hormone (GnRH)-induced gonadotropin secretion and regulates the pubertal activation of GnRH nuerons. A polymorphism in the terminal exon of this mRNA results in two protein isoforms. An adenosine present at the polymorphic site represents the third position in a stop codon. When the adenosine is absent, a downstream stop codon is utilized and the encoded protein extends for an additional seven amino acid residues.

[provided by RefSeq, Mar 2012]

Function Metastasis suppressor protein in malignant melanomas and in some breast cancers. May regulate

events downstream of cell-matrix adhesion, perhaps involving cytoskeletal reorganization. Generates a C-terminally amidated peptide, metastin which functions as the endogenous ligand of the G-protein coupled receptor GPR54. Activation of the receptor inhibits cell proliferation and cell migration, key characteristics of tumor metastasis. Kp-10 is a decapeptide derived from the primary translation product, isolated in conditioned medium of first trimester trophoblast. Kp-10, but not other kisspeptins, increased intracellular Ca(2+) levels in isolated first trimester trophoblasts. Kp-10 is a

paracrine/endocrine regulator in fine-tuning trophoblast invasion generated by the trophoblast itself. The receptor is also essential for normal gonadotropin-released hormone physiology and for puberty. The hypothalamic KiSS1/GPR54 system is a pivotal factor in central regulation of the gonadotropic axis


at puberty and in adulthood. [UniProt]

Calculated Mw 15 kDa

PTM Processed by MMP2 and MMP9. [UniProt]


Cellular Localization Secreted. [UniProt]

Images



ARG59533 anti-Kisspeptin antibody IHC-P image

Immunohistochemistry: Paraformaldehyde-fixed and paraffinembedded Human testis section blocked with 3% BSA for 0.5 hour at RT. Antigen Retrieval: Heat mediation was performed in Citrate buffer (pH 6.0). Samples were stained with ARG59533 anti-Kisspeptin antibody at 1:25 for 1 hours at 37°C.



ARG59533 anti-Kisspeptin antibody WB image

Western blot: 20 μg of Human skeletal muscle lysate stained with ARG59533 anti-Kisspeptin antibody at 1:2000 dilution.

Human skeletal muscle