

Product datasheet

info@arigobio.com

ARG59726 anti-SUMO4 (M55V) antibody

Package: 100 μl Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes SUMO4 (M55V)

Tested Reactivity Hu

Tested Application IHC-P, WB

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name SUMO4 (M55V)

Species Human

Immunogen KLH-conjugated synthetic peptide of Human SUMO4. (CEPRGLS(V)KQIRFRFG)

Conjugation Un-conjugated

Alternate Names SUMO-4; IDDM5; SMT3H4; dJ281H8.4; Small ubiquitin-related modifier 4; Small ubiquitin-like protein 4

Application Instructions

Application table	Application	Dilution
	IHC-P	1:50 - 1:100
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HepG2	

Properties

_	
Form	Liguid

Purification This antibody is affinity purified using peptides CEPRGLS(V)KQIRFRFG (positive selection) and

CEPRGLS(M)KQIRFRFG (negative selection).

Buffer PBS and 0.09% (W/V) Sodium azide.

Preservative 0.09% (W/V) Sodium azide.

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol

Gene Full Name small ubiquitin-like modifier 4

SUMO4

Background This gene is a member of the SUMO gene family. This family of genes encode small ubiquitin-related

modifiers that are attached to proteins and control the target proteins' subcellular localization, stability, or activity. The protein described in this record is located in the cytoplasm and specifically modifies IKBA, leading to negative regulation of NF-kappa-B-dependent transcription of the IL12B gene. A specific polymorphism in this SUMO gene, which leads to the M55V substitution, has been associated

with type I diabetes. The RefSeq contains this polymorphism. [provided by RefSeq, Jul 2008]

Function Ubiquitin-like protein which can be covalently attached to target lysines as a monomer. Does not seem

to be involved in protein degradation and may modulate protein subcellular localization, stability or activity. Upon oxidative stress, conjugates to various anti-oxidant enzymes, chaperones, and stress defense proteins. May also conjugate to NFKBIA, TFAP2A and FOS, negatively regulating their transcriptional activity, and to NR3C1, positively regulating its transcriptional activity. Covalent attachment to its substrates requires prior activation by the E1 complex SAE1-SAE2 and linkage to the

E2 enzyme UBE2I. [UniProt]


Calculated Mw 11 kDa

PTM In contrast to SUMO1, SUMO2 and SUMO3, seems to be insensitive to sentrin-specific proteases due to

the presence of Pro-90. This may impair processing to mature form and conjugation to substrates.


[UniProt]

Images



ARG59726 anti-SUMO4 (M55V) antibody IHC-P image

Immunohistochemistry: Formalin-fixed and paraffin-embedded Human hepatocarcinoma stained with ARG59726 anti-SUMO4 (M55V) antibody.



HepG2

ARG59726 anti-SUMO4 (M55V) antibody WB image

Western blot: 35 μg of HepG2 cell lysate stained with ARG59726 anti-SUMO4 (M55V) antibody.