

## ARG62472 anti-DP1 antibody

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes DP1
Tested Reactivity	Hu, Ms, Rat
Tested Application	IP, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	DP1
Species	Human
Immunogen	Synthetic peptide from the N terminus of human DP-1 protein.
Epitope	N-terminal
Conjugation	Un-conjugated
Alternate Names	DRTF1-polypeptide 1; DP1; DRTF1; E2F dimerization partner 1; Transcription factor Dp-1; Dp-1

### Application Instructions

Application table	Application	Dilution
	IP	1:2000
	WB	1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	A431 or Raji cells	

### Properties

Form	Liquid
Purification	Purified Antibody
Buffer	1X PBS and 0.1% Sodium azide
Preservative	0.1% Sodium azide
Concentration	0.2 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Database links	<a href="#">GeneID: 21781 Mouse</a> <a href="#">GeneID: 7027 Human</a> <a href="#">Swiss-port # Q08639 Mouse</a> <a href="#">Swiss-port # Q14186 Human</a>
Gene Symbol	TFDP1
Gene Full Name	transcription factor Dp-1
Background	This gene encodes a member of a family of transcription factors that heterodimerize with E2F proteins to enhance their DNA-binding activity and promote transcription from E2F target genes. The encoded protein functions as part of this complex to control the transcriptional activity of numerous genes involved in cell cycle progression from G1 to S phase. Alternative splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 1, 15, and X.[provided by RefSeq, Jan 2009]
Function	Can stimulate E2F-dependent transcription. Binds DNA cooperatively with E2F family members through the E2 recognition site, 5'-TTTC[CG]CGC-3', found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication. The E2F1:DP complex appears to mediate both cell proliferation and apoptosis. Blocks adipocyte differentiation by repressing CEBPA binding to its target gene promoters (PubMed:20176812). [UniProt]
Research Area	Developmental Biology antibody; Gene Regulation antibody; Signaling Transduction antibody
Calculated Mw	45 kDa
PTM	Phosphorylation by E2F1-bound cyclin A-CDK2, in the S phase, inhibits E2F-mediated DNA binding and transactivation. Ubiquitinated by the BCR(KBTBD5) complex, leading to its subsequent degradation.