

Product datasheet

info@arigobio.com

ARG62541 anti-MCP5 / Mast cell chymase antibody [CC1]

Package: 100 μl Store at: -20°C

Summary

Product Description Mouse Monoclonal antibody [CC1] recognizes MCP5 / Mast cell chymase

Tested Reactivity Hu, Rat, Pig

Tested Application FACS, IHC-Fr, IHC-P, WB

Host Mouse

Clonality Monoclonal

Clone CC1

Isotype IgG1

Target Name MCP5 / Mast cell chymase

Species Human

Immunogen Purified human skin Chymase

Conjugation Un-conjugated

Alternate Names Chymase; Alpha-chymase; Mast cell protease I; chymase; CYH; EC 3.4.21.39; MCT1

Application Instructions

Application Note FACS: 1-2µg for 106 cells

IHC: 1/100-1/200

* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations

should be determined by the scientist.

Properties

Form Liquid

Purification Immunogen affinity purified

Buffer PBS, 1% BSA and 0.05% Sodium azide

Preservative 0.05% Sodium azide

Stabilizer 1% BSA

Concentration 0.2 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links GenelD: 1215 Human

GeneID: 25627 Rat

Swiss-port # P23946 Human

Swiss-port # P50339 Rat

Gene Symbol CMA1

Gene Full Name chymase 1, mast cell

Background This gene encodes a chymotryptic serine proteinase that belongs to the peptidase family S1. It is

expressed in mast cells and is thought to function in the degradation of the extracellular matrix, the regulation of submucosal gland secretion, and the generation of vasoactive peptides. In the heart and blood vessels, this protein, rather than angiotensin converting enzyme, is largely responsible for converting angiotensin I to the vasoactive peptide angiotensin II. Alternative splicing results in multiple

variants. [provided by RefSeq, Apr 2015]

Function Major secreted protease of mast cells with suspected roles in vasoactive peptide generation,

extracellular matrix degradation, and regulation of gland secretion. [UniProt]

Research Area Cell Biology and Cellular Response antibody; Immune System antibody; Signaling Transduction antibody

Calculated Mw 27 kDa

Cellular Localization Secreted, Cytoplasmic granule