

Product datasheet

info@arigobio.com

ARG62908 anti-CD62L / L-Selectin antibody [IVA94]

Package: 100 μg Store at: -20°C

Summary

Product Description Mouse Monoclonal antibody [IVA94] recognizes CD62L / L-Selectin

Tested Reactivity Bov

Tested Application FACS, ICC/IF, IP

Specificity The clone IVA94 reacts with CD62L antigen (bovine). CD62L (L-selectin) is a 74-95 kDa single chain type I

glycoprotein expressed on most peripheral blood B lymphocytes, T lymphocytes, monocytes and granulocytes; it is also present on a subset of NK cells and certain hematopoietic malignant cells.

Host Mouse

Clonality Monoclonal

Clone IVA94

Isotype IgG1

Target Name CD62L / L-Selectin

Immunogen Bovine leukocytes.

Conjugation Un-conjugated

Alternate Names Leukocyte surface antigen Leu-8; Leukocyte adhesion molecule 1; CD antigen CD62L; PLNHR; LSEL;

CD62L; Leukocyte-endothelial cell adhesion molecule 1; L-selectin; LAM1; LNHR; TQ1; CD62 antigen-like

family member L; gp90-MEL; Lymph node homing receptor; LYAM1; LECAM1; LEU8; LAM-1

Application Instructions

Application table	Application	Dilution
	FACS	1 - 4 μg/ml
	ICC/IF	Assay-dependent
	IP	Assay-dependent
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form Liquid

Purification Purified from ascites by protein-A affinity chromatography.

Purity > 95% (by SDS-PAGE)

Buffer PBS (pH 7.4) and 15 mM Sodium azide

Preservative 15 mM Sodium azide

Concentration 1 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links GeneID: 281485 Bovine

Swiss-port # P98131 Bovine

Gene Symbol SELL

Gene Full Name selectin L

Background This gene encodes a cell surface adhesion molecule that belongs to a family of adhesion/homing

receptors. The encoded protein contains a C-type lectin-like domain, a calcium-binding epidermal growth factor-like domain, and two short complement-like repeats. The gene product is required for binding and subsequent rolling of leucocytes on endothelial cells, facilitating their migration into secondary lymphoid organs and inflammation sites. Single-nucleotide polymorphisms in this gene have been associated with various diseases including immunoglobulin A nephropathy. Alternatively spliced

transcript variants have been found for this gene. [provided by RefSeq, Oct 2009]

Function Cell surface adhesion protein. Mediates the adherence of lymphocytes to endothelial cells of high

endothelial venules in peripheral lymph nodes. Promotes initial tethering and rolling of leukocytes in


endothelia (By similarity). [UniProt]

Research Area Cell Biology and Cellular Response antibody; Developmental Biology antibody; Immune System

antibody; Signaling Transduction antibody


Calculated Mw 42 kDa

Images




ARG62908 anti-CD62L / L-Selectin antibody [IVA94] ICC/IF image

Immunofluorescence: Acetone / methanol-fixed Bovine peripheral blood (lymphocyte-enriched suspension) stained with ARG62908 anti-CD62L / L-Selectin antibody [IVA94] (green), DAPI (blue) for nuclear staining.



ARG62908 anti-CD62L / L-Selectin antibody [IVA94] FACS image

Flow Cytometry: Separation of bovine CD62L positive lymphocytes (red-filled) from CD62L negative lymphocytes (black-dashed). Bovine peripheral whole blood stained with ARG62908 anti-CD62L / L-Selectin antibody [IVA94] at 1 $\mu g/ml$ dilution, followed by APC-conjugated Goat anti-Mouse antibody.



ARG62908 anti-CD62L / L-Selectin antibody [IVA94] FACS image

Flow Cytometry: Bovine peripheral whole blood stained with ARG62908 anti-CD62L / L-Selectin antibody [IVA94] at 1 $\mu g/ml$ dilution, followed by APC-conjugated Goat anti-Mouse antibody.