

**ARG63304**  
**anti-RNF35 / TRIM40 antibody**

Package: 100 µg

Store at: -20°C

### Summary

Product Description	Goat Polyclonal antibody recognizes RNF35 / TRIM40
Tested Reactivity	Hu
Tested Application	WB
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	RNF35 / TRIM40
Species	Human
Immunogen	C-GVSELLQPPQKL
Conjugation	Un-conjugated
Alternate Names	RING finger protein 35; RNF35; Probable E3 NEDD8-protein ligase; Tripartite motif-containing protein 40

### Application Instructions

Application table	Application	Dilution
	WB	0.5 - 2 µg/ml
Application Note	WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---

Database links [GeneID: 135644 Human](#)

[Swiss-port # Q6P9F5 Human](#)


**Background** This gene encodes a member of the tripartite motif (TRIM) protein family. The encoded protein may play a role as a negative regulator against inflammation and carcinogenesis in the gastrointestinal tract. Alternatively spliced transcript variants that encode different protein isoforms have been described. [provided by RefSeq, Feb 2014]

**Research Area** Cell Biology and Cellular Response antibody

**Calculated Mw** 29 kDa

## Images

---



ARG63304 anti-RNF35 / TRIM40 antibody WB image

Western Blot: Human Placenta lysate (35 µg protein in RIPA buffer) stained with ARG63304 anti-RNF35 / TRIM40 antibody at 0.5 µg/ml dilution.