

ARG63308 anti-TRIM35 / MAIR antibody

Package: 100 μg Store at: -20°C

Summary

Product Description	Goat Polyclonal antibody recognizes TRIM35 / MAIR
Tested Reactivity	Hu
Tested Application	WB
Host	Goat
Clonality	Polyclonal
Isotype	lgG
Target Name	TRIM35 / MAIR
Species	Human
Immunogen	ERSPDVSPGPSRS-C
Conjugation	Un-conjugated
Alternate Names	Hemopoietic lineage switch protein 5; Tripartite motif-containing protein 35; HLS5; MAIR

Application Instructions

Application table	Application	Dilution
	WB	1 - 3 μg/ml
Application Note	WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid	
Purification	Purified from goat serum by antigen affinity chromatography.	
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.	
Preservative	0.02% Sodium azide	
Stabilizer	0.5% BSA	
Concentration	0.5 mg/ml	
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.	
Note	For laboratory research only, not for drug, diagnostic or other use.	

Bioinformation

Database links	GenelD: 23087 Human
	Swiss-port # Q9UPQ4 Human
Background	The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The function of this protein has not been identified. [provided by RefSeq, Jul 2008]
Research Area	Cell Biology and Cellular Response antibody; Gene Regulation antibody
Calculated Mw	57 kDa

Images

250kDa 150kDa 100kDa 75kDa	ARG63308 anti-TRIM35 / MAIR antibody WB image Western Blot: K562 lysate (35 μg protein in RIPA buffer) stained with ARG63308 anti-TRIM35 / MAIR antibody at 1 μg/ml dilution.
50kDa	
37kDa	
25kDa	
20kDa	
20104	
15kDa	