

ARG63370 anti-TCF19 / SC1 antibody

Package: 100 µg
Store at: -20°C

Summary

Product Description	Goat Polyclonal antibody recognizes TCF19 / SC1
Tested Reactivity	Hu
Tested Application	WB
Specificity	Immunizing peptide was designed based on NP_009040.1, but this version has become NP_009040.2 and shorter, thus no longer matching with the peptide.
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	TCF19 / SC1
Species	Human
Immunogen	C-RSTAKAPSDTPAHE
Conjugation	Un-conjugated
Alternate Names	TCF-19; Transcription factor 19; SC1; Transcription factor SC1

Application Instructions

Application table	Application	Dilution
	WB	1 - 2 µg/ml
Application Note	WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 6941 Human Swiss-port # Q9Y242 Human
Background	This gene encodes a protein that belongs to the Plant Homeo Domain finger family of transcription factors. The encoded protein is thought to function during the G1/S transition in the cell cycle. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Sep 2013]
Research Area	Developmental Biology antibody; Gene Regulation antibody; Signaling Transduction antibody
Calculated Mw	37 kDa

Images



ARG63370 anti-TCF19 / SC1 antibody WB image

Western Blot: 293 lysate (RIPA buffer, 30µg total protein per lane) stained with ARG63370 anti-TCF19 antibody at 1.5 µg/ml dilution.