

Product datasheet

info@arigobio.com

ARG63649 anti-BAZ2B antibody

Package: 100 μg Store at: -20°C

Summary

Product Description Goat Polyclonal antibody recognizes BAZ2B

Tested Reactivity Hu

Predict Reactivity Ms, Rat, Cow

Tested Application IHC-P
Host Goat

Clonality Polyclonal

Isotype IgG

Target Name BAZ2B
Species Human

Immunogen C-YFEKKWTDTFKVS

Conjugation Un-conjugated

Alternate Names WALp4; hWALp4; Bromodomain adjacent to zinc finger domain protein 2B

Application Instructions

Application table	Application	Dilution
	IHC-P	1 - 2 μg/ml
Application Note	IHC-P: Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0). * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form Liquid

Purification Purified from goat serum by antigen affinity chromatography.

Buffer Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.

Preservative 0.02% Sodium azide

Stabilizer 0.5% BSA

Concentration 0.5 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

www.arigobio.com arigo.nuts about antibodies 1/2

Bioinformation

Database links GeneID: 29994 Human

Swiss-port # Q9UIF8 Human

Gene Symbol BAZ2B


Gene Full Name bromodomain adjacent to zinc finger domain, 2B

Function May play a role in transcriptional regulation interacting with ISWI. [UniProt]

Research Area Gene Regulation antibody

Calculated Mw 240 kDa

Images



ARG63649 anti-BAZ2B antibody IHC-P image

Immunohistochemistry: paraffin embedded Human Small Intestine. (Steamed antigen retrieval with citrate buffer pH 6) stained with ARG63649 anti-BAZ2B antibody at 5 $\mu g/ml$ dilution followed by AP-staining.