

**ARG63837**  
anti-VPS41 antibodyPackage: 100 µg  
Store at: -20°C

### Summary

Product Description	Goat Polyclonal antibody recognizes VPS41
Tested Reactivity	Hu
Predict Reactivity	Ms, Rat, Cow, Dog
Tested Application	IHC-P
Specificity	This antibody is expected to recognise both reported isoforms (NP_055211.2 and NP_542198.2)
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	VPS41
Species	Human
Immunogen	C-DHLKKDSQNKTLK
Conjugation	Un-conjugated
Alternate Names	HVSP41; Vacuolar protein sorting-associated protein 41 homolog; S53; HVPS41; hVps41p

### Application Instructions

Application table	Application	Dilution
	IHC-P	3 - 6 µg/ml

**Application Note** IHC-P: Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0).  
\* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

### Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

Database links

[GeneID: 27072 Human](#)

[Swiss-port # P49754 Human](#)

Background

Vesicle mediated protein sorting plays an important role in segregation of intracellular molecules into distinct organelles. Genetic studies in yeast have identified more than 40 vacuolar protein sorting (VPS) genes involved in vesicle transport to vacuoles. This gene encodes the human ortholog of yeast Vps41 protein which is also conserved in Drosophila, tomato, and Arabidopsis. Expression studies in yeast and human indicate that this protein may be involved in the formation and fusion of transport vesicles from the Golgi. Several transcript variants encoding different isoforms have been described for this gene, however, the full-length nature of not all is known. [provided by RefSeq, Jul 2008]


Research Area

Signaling Transduction antibody

Calculated Mw


99 kDa

## Images




ARG63837 anti-VPS41 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human placenta tissue. Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0). The tissue section was stained with ARG63837 anti-VPS41 antibody at 3.75 µg/ml dilution followed by AP-staining.



ARG63837 anti-VPS41 antibody IHC-P image

Immunohistochemistry: paraffin embedded Human Kidney. (Steamed antigen retrieval with citrate buffer pH 6) stained with ARG63837 anti-VPS41 antibody at 4 µg/ml dilution followed by HRP-staining. Similar results were obtained after antigen retrieval at pH9.



#### ARG63837 anti-VPS41 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human prostate tissue.  
Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0). The tissue section was stained with ARG63837 anti-VPS41 antibody at 3.75 µg/ml dilution followed by AP-staining.