

ARG64026 anti-BMPR1A antibody

Package: 100 µg
Store at: -20°C

Summary

| | |
|---------------------|--|
| Product Description | Goat Polyclonal antibody recognizes BMPR1A |
| Tested Reactivity | Hu |
| Predict Reactivity | Dog |
| Tested Application | WB |
| Host | Goat |
| Clonality | Polyclonal |
| Isotype | IgG |
| Target Name | BMPR1A |
| Species | Human |
| Immunogen | C-KSDSDQKKSEN |
| Conjugation | Un-conjugated |
| Alternate Names | CD292; CD antigen CD292; ALK3; ACVRLK3; ALK-3; EC 2.7.11.30; Activin receptor-like kinase 3; 10q23del; Serine/threonine-protein kinase receptor R5; BMPR-1A; BMP type-1A receptor; SKR5; Bone morphogenetic protein receptor type-1A |

Application Instructions

| Application table | Application | Dilution |
|-------------------|--|-------------|
| | WB | 1 - 3 µg/ml |
| Application Note | WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. | |

Properties

| | |
|---------------------|--|
| Form | Liquid |
| Purification | Purified from goat serum by antigen affinity chromatography. |
| Buffer | Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA. |
| Preservative | 0.02% Sodium azide |
| Stabilizer | 0.5% BSA |
| Concentration | 0.5 mg/ml |
| Storage instruction | For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. |

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

| | |
|----------------|--|
| Database links | GeneID: 657 Human Swiss-port # P36894 Human |
| Background | The bone morphogenetic protein (BMP) receptors are a family of transmembrane serine/threonine kinases that include the type I receptors BMPR1A and BMPR1B and the type II receptor BMPR2. These receptors are also closely related to the activin receptors, ACVR1 and ACVR2. The ligands of these receptors are members of the TGF-beta superfamily. TGF-betas and activins transduce their signals through the formation of heteromeric complexes with 2 different types of serine (threonine) kinase receptors: type I receptors of about 50-55 kD and type II receptors of about 70-80 kD. Type II receptors bind ligands in the absence of type I receptors, but they require their respective type I receptors for signaling, whereas type I receptors require their respective type II receptors for ligand binding. [provided by RefSeq, Jul 2008] |
| Research Area | Cell Biology and Cellular Response antibody; Developmental Biology antibody; Metabolism antibody; Signaling Transduction antibody |
| Calculated Mw | 60 kDa |

Images



ARG64026 anti-BMPR1A antibody WB image

Western Blot: HeLa cell lysate (35 µg protein in RIPA buffer) stained with ARG64026 anti-BMPR1A antibody at 1 µg/ml dilution.