

#### ARG64035 anti-APBB1 / FE65 antibody

Package: 100 μg Store at: -20°C

# Summary

Product Description	Goat Polyclonal antibody recognizes APBB1 / FE65
Tested Reactivity	Ms
Predict Reactivity	Hu
Tested Application	WB
Specificity	This antibody is expected to recognise both reported isoforms (NP_001155.1 and NP_663722.1).
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	APBB1 / FE65
Species	Human
Immunogen	C-GSLKPKRLGAHTP
Conjugation	Un-conjugated
Alternate Names	FE65; Amyloid beta A4 precursor protein-binding family B member 1; Protein Fe65; RIR; MGC:9072

#### **Application Instructions**

Application table	Application	Dilution		
	WB	0.1 - 0.3 μg/ml		
Application Note	WB: Recommend incubate at RT for 1h.			
		* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.		

## Properties

Form	Liquid	
Purification	Purified from goat serum by antigen affinity chromatography.	
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.	
Preservative	0.02% Sodium azide	
Stabilizer	0.5% BSA	
Concentration	0.5 mg/ml	
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.	

### **Bioinformation**

Database links	GeneID: 11785 Mouse	
	Swiss-port # Q9QXJ1 Mouse	
Background	The protein encoded by this gene is a member of the Fe65 protein family. It is an adaptor protein localized in the nucleus. It interacts with the Alzheimer's disease amyloid precursor protein (APP), transcription factor CP2/LSF/LBP1 and the low-density lipoprotein receptor-related protein. APP functions as a cytosolic anchoring site that can prevent the gene product's nuclear translocation. This encoded protein could play an important role in the pathogenesis of Alzheimer's disease. It is thought to regulate transcription. Also it is observed to block cell cycle progression by downregulating thymidylate synthase expression. Multiple alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Mar 2012]	
Research Area	Neuroscience antibody; Signaling Transduction antibody	
Calculated Mw	77 kDa	
PTM	Phosphorylation at Ser-610 by SGK1 promotes its localization to the nucleus (By similarity). Phosphorylated following nuclear translocation. Phosphorylation at Tyr-547 by ABL1 enhances transcriptional activation activity and reduces the affinity for RASD1/DEXRAS1.	

## Images

250kDa 150kDa	ARG64035 anti-APBB1 / FE65 antibody WB image
- 100kDa 75kDa 50kDa	Western Blot: NIH/3T3 lysate (35 $\mu g$ protein in RIPA buffer) stained with ARG64035 anti-APBB1 / FE65 antibody at 0.1 $\mu g/ml$ dilution.
37kDa	
25kDa	
20kDa	
15kDa	
10kDa	