

ARG64464 anti-TEB4 / MARCH VI antibody

Package: 100 µg
Store at: -20°C

Summary

Product Description	Goat Polyclonal antibody recognizes TEB4 / MARCH VI
Tested Reactivity	Hu
Predict Reactivity	Ms
Tested Application	WB
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	TEB4 / MARCH VI
Species	Human
Immunogen	C-QGRRVIFQKVKE
Conjugation	Un-conjugated
Alternate Names	RNF176; EC 6.3.2.-; Membrane-associated RING finger protein 6; TEB4; RING finger protein 176; DOA10; E3 ubiquitin-protein ligase MARCH6; Doa10 homolog; MARCH-VI; Membrane-associated RING-CH protein VI; Protein TEB-4

Application Instructions

Application table	Application	Dilution
	WB	0.2 - 0.5 µg/ml
Application Note	WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties


Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 10299 Human Swiss-port # O60337 Human
Background	This gene encodes a member of a family of membrane-associated E3 ubiquitin ligases containing RING-CH-type zinc finger motifs. Ubiquitination of type II deiodinase by the encoded protein is an important regulatory step in thyroid hormone signalling. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Jul 2012]
Research Area	Cell Biology and Cellular Response antibody
Calculated Mw	103 kDa
PTM	Auto-ubiquitinated, which results in proteasomal degradation.

Images



ARG64464 anti-TEB4 / MARCH VI antibody WB image

Western blot: Human Lung lysate (35 µg protein in RIPA buffer) stained with ARG64464 anti-TEB4 / MARCH VI antibody at 0.2 µg/ml dilution.