

Summary

ARG65533 anti-CD45R / B220 antibody [RA3-6B2] (low endotoxin)

Package: 100 μg Store at: -20°C

,	
Product Description	Azide free and low endotoxin Rat Monoclonal antibody [RA3-6B2] recognizes CD45R
Tested Reactivity	Hu, Ms, Cat
Tested Application	FACS, FuncSt, ICC/IF, IHC-Fr, IHC-P, IP
Specificity	The clone RA3-6B2 recognizes an epitope on CD45R, which is expressed at all developmental stages of B cells, including activated B cells, but also on subsets of NK and T cells. T cells detected by this antibody are supposed to be in activated state.
Host	Rat
Clonality	Monoclonal
Clone	RA3-6B2
Isotype	lgG2a
Target Name	CD45R / B220
Immunogen	Abelson murine leukemia virus-induced pre-B tumor cells
Conjugation	Un-conjugated
Alternate Names	LY5; GP180; Receptor-type tyrosine-protein phosphatase C; CD45; L-CA; CD antigen CD45; Leukocyte common antigen; CD45R; LCA; T200; EC 3.1.3.48; B220

Application Instructions

Application table	Application	Dilution
	Application	Dilution
	FACS	1 - 4 µg/ml
	FuncSt	Assay-dependent
	ICC/IF	Assay-dependent
	IHC-Fr	Assay-dependent
	IHC-P	Assay-dependent
	IP	Assay-dependent
Application Note		dulation of B cell responses. recommended starting dilutions and the optimal dilutions or concentrations by the scientist.

Properties

Form	Liquid
Purification	Purification with Protein G.
Purification Note	0.2 μm filter sterilized. Endotoxin level is 95% (by SDS-PAGE)

Buffer	PBS (pH 7.4)
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GenelD: 19264 Mouse
	GeneID: 5788 Human
	Swiss-port # P08575 Human
Background	CD45 is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitosis, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus is classified as a receptor type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported. [provided by RefSeq, Jun 2012]
Function	CD45: Protein tyrosine-protein phosphatase required for T-cell activation through the antigen receptor. Acts as a positive regulator of T-cell coactivation upon binding to DPP4. The first PTPase domain has enzymatic activity, while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation, recruits and dephosphorylates SKAP1 and FYN. Dephosphorylates LYN, and thereby modulates LYN activity.
Highlight	 (Microbial infection) Acts as a receptor for human cytomegalovirus protein UL11 and mediates binding of UL11 to T-cells, leading to reduced induction of tyrosine phosphorylation of multiple signaling proteins upon T-cell receptor stimulation and impaired T-cell proliferation. [UniProt] Related products: <u>CD45R antibodies</u>; <u>CD45R Duos / Panels</u>; <u>Anti-Rat IgG secondary antibodies</u>; Related news: <u>Exploring Antiviral Immune Response</u>
Research Area	Developmental Biology antibody; Immune System antibody; Neuroscience antibody; Signaling Transduction antibody; Mouse Inflammatory Cell Marker antibody; B Cell Marker antibody
Calculated Mw	147 kDa
PTM	Heavily N- and O-glycosylated.